

ISSN 1342-2804

Research Reports on Mathematical and Computing Sciences

A Lagrangian-DNN Relaxation: a Fast Method
for Computing Tight Lower Bounds for a Class
of Quadratic Optimization Problems

Sunyoung Kim, Masakazu Kojima
and Kim-Chuan Toh

October 2013, B-472

Department of
Mathematical and
Computing Sciences
Tokyo Institute of Technology

SERIES **B:** Operations Research

B-472 A Lagrangian-DNN Relaxation: a Fast Method for Computing Tight Lower Bounds for a Class of Quadratic Optimization Problems

Sunyoung Kim^{*}, Masakazu Kojima[†] and Kim-Chuan Toh[‡]

October 2013

Abstract.

We propose an efficient computational method for linearly constrained quadratic optimization problems (QOPs) with complementarity constraints based on their Lagrangian and doubly nonnegative (DNN) relaxation and first-order algorithms. The simplified Lagrangian-CPP relaxation of such QOPs proposed by Arima, Kim, and Kojima in 2012 takes one of the simplest forms, an unconstrained conic linear optimization problem with a single Lagrangian parameter in a completely positive (CPP) matrix variable with its upper-left element fixed to 1. Replacing the CPP matrix variable by a DNN matrix variable, we derive the Lagrangian-DNN relaxation, and establish the equivalence between the optimal value of the DNN relaxation of the original QOP and that of the Lagrangian-DNN relaxation. We then propose an efficient numerical method for the Lagrangian-DNN relaxation using a bisection method combined with the proximal alternating direction multiplier and the accelerated proximal gradient methods. Numerical results on binary QOPs, quadratic multiple knapsack problems, maximum stable set problems, and quadratic assignment problems illustrate the superior performance of the proposed method for attaining tight lower bounds in shorter computational time.

Keywords: Linearly constrained quadratic optimization problems with complementarity constraints, the Lagrangian-conic relaxation, Bisection method, Iterative solver.

AMS Classification. 90C20, 90C25, 90C26.

^{*} Department of Mathematics, Ewha W. University, 11-1 Dahyun-dong, Sudaemoongu, Seoul 120-750 Korea. The research was supported by NRF 2012-R1A1A2-038982 (skim@ewha.ac.kr).

[†] Research and Development Initiative & JST CREST, Chuo University, 1-13-27, Kasuga, Bunkyo-ku, Tokyo 112-8551 Japan. This research was supported by the Japan Science and Technology Agency (JST), the Core Research of Evolutionary Science and Technology (CREST) research project (kojima@is.titech.ac.jp).

[‡] Department of Mathematics, National University of Singapore, 10 Lower Kent Ridge Road, Singapore 119076. Research supported in part by Ministry of Education Academic Research Fund R-146-000-168-112 (mattohkc@nus.edu.sg).

1 Introduction

We consider a linearly constrained quadratic optimization problem (QOP) with complementarity constraints:

$$\text{minimize } \left\{ \mathbf{u}^T \mathbf{Q} \mathbf{u} + 2\mathbf{c}^T \mathbf{u} \mid \begin{array}{l} \mathbf{u} \in \mathbb{R}_+^m, \mathbf{A}\mathbf{u} + \mathbf{b} = \mathbf{0}, \\ u_i u_j = 0 \ ((i, j) \in \mathcal{E}) \end{array} \right\} \quad (1)$$

where $\mathbf{A} \in \mathbb{R}^{q \times m}$, $\mathbf{b} \in \mathbb{R}^q$, $\mathbf{c} \in \mathbb{R}^m$ and $\mathcal{E} \subset \{(i, j) : 1 \leq i < j \leq m\}$ are given data. Noting that the binary constraint $u_i(1 - u_i) = 0$ can be converted to a complementarity constraint $u_i v_i = 0$ with a slack variable $v_i = 1 - u_i \geq 0$, thus QOP (1) can model nonconvex quadratic problems with linear, binary and complementarity constraints. We assume that the linear constraint set $\{\mathbf{u} \in \mathbb{R}_+^m : \mathbf{A}\mathbf{u} + \mathbf{b} = \mathbf{0}\}$ is bounded. The QOP model (1) satisfying this assumption includes various combinatorial optimization problems, for instance, the binary integer quadratic problem, the maximum stable set problem, the quadratic multiple knapsack problem, and the quadratic assignment problem [9, 23, 25].

The completely positive programming (CPP) relaxation of linearly constrained QOPs in binary and continuous variables by Burer [8] has gained considerable attention since 2009. Extending his work, theoretical results for a more general class of QOPs were established by Eichfelder and Povh [11, 13] and by Arima, Kim and Kojima [1]. They showed that the exact optimal values of QOPs in their classes coincide with the optimal values of their CPP relaxation problems.

Such CPP relaxations are numerically intractable since the problem of determining whether a given matrix lies in the completely positive cone and the copositive cone is a co-NP-complete problem as shown in [21]. Replacing the completely positive cone by doubly nonnegative (DNN) cone and solving the resulting problem by a primal-dual interior-point method has been a popular approach [17, 31]. The computational cost of this approach, however, is very expensive as the number of nonnegative constraints for the variables grows rapidly with the size of the problem.

Recently, Arima, Kim and Kojima [2] introduced the simplified Lagrangian-CPP relaxation of a linearly constrained QOP in continuous and binary variables with a single parameter $\lambda \in \mathbb{R}$. It was derived by reducing the original QOP to an equivalent QOP with a single quadratic equality constraint in nonnegative variables, and applying the Lagrangian relaxation to the resulting QOP. As a result, an unconstrained QOP with a Lagrangian multiplier $\lambda \in \mathbb{R}$ in nonnegative variables was obtained. From the computational point of view, this Lagrangian relaxation is one of the simplest forms to handle with a solution method. Applying the CPP relaxation to the unconstrained QOP with λ lead to the Lagrangian-CPP relaxation. It was shown in [2] that the optimal values of the Lagrangian relaxation as well as its CPP relaxation monotonically converge to the exact optimal value of the original QOP as λ tends to ∞ .

The main goals of this paper are twofold. First, we propose an efficient and effective numerical method for the QOP model (1) by extending the framework of the Lagrangian-CPP relaxation to the one including the Lagrangian-DNN relaxation of (1). Second,

generalizing the brief discussion on such an extension in [2], we present a theoretical framework for the Lagrangian-conic relaxation of (1) that covers both Lagrangian-CPP and Lagrangian-DNN relaxations. We use the primal-dual pair of the Lagrangian-DNN relaxation with a sufficiently large $\lambda \in \mathbb{R}$ to compute a tight lower bound for the optimal value of (1). The main features of the Lagrangian-DNN relaxation are:

- The primal is an unconstrained DNN problem with a matrix variable whose upper-left corner element is fixed to 1. Thus, its dual becomes a simple problem with just a single variable.
- The primal DNN problem is strictly feasible, *i.e.*, the primal feasible region intersects with the interior of the DNN cone.
- A common optimal objective value, shared by the primal-dual pair with a parameter $\lambda > 0$, monotonically converges to the optimal objective value of the DNN relaxation of (1). Hence, a lower bound with almost the same quality as the one obtained from the DNN relaxation of (1) can be computed for the optimal objective value of (1) via the simple Lagrangian-DNN relaxation with a sufficiently large $\lambda > 0$.

The computational efficiency for solving the Lagrangian-DNN relaxation can be expected from the first and second features mentioned above. If a primal-dual interior-point method [7, 15, 28, 29] for SDPs is used to solve the Lagrangian-DNN relaxation, then the inequality constraints induced from the nonnegativity of all elements of the DNN matrix variable may incur prohibitive computational burden. More precisely, if the size of the DNN matrix is n , then the number of inequalities to be added in the SDP problem amounts to $n(n-1)/2$. Thus, the conversion of a DNN problem into a standard SDP is computationally inefficient when n is large. To avoid such inefficiency of using a primal-dual interior-point method, the numerical method proposed in this paper employs *first-order algorithms* [4] for the Lagrangian-DNN relaxation without converting it into a standard SDP. The first and second features mentioned previously considerably increase the efficiency and numerical stability of the first-order algorithms, respectively.

The numerical results in Section 5 show that the Lagrangian-DNN relaxation provides tighter lower bounds more efficiently than the DNN relaxation of the QOP (1). When the proposed method is experimented on the test problems such as the binary integer quadratic problem, the maximum stable set problem, the quadratic multiple knapsack problem, and the quadratic assignment problem, the quality of the lower bounds obtained from the proposed method is tight, compared to the known optimal values. As mentioned in the third main feature, a sufficiently large λ results in a tight lower bound. The proposed method can also solve the problems efficiently, in particular, it obtains the lower bound for the quadratic assignment problem much faster than SDPNAL, which is an advanced large scale SDP solver [32], applied to the DNN relaxation of the problem.

In Section 2, we list notation and symbols used throughout the paper. The QOP (1) is extended to a general QOP model, from which effective conic and Lagrangian-conic

relaxations are derived. We also describe sufficient conditions for them to attain the same optimal value. In Section 3, the conic and Lagrangian-conic relaxations and their relations are discussed. In particular, the Lagrangian-DNN relaxation, a special case of the Lagrangian-conic relaxations, is used to obtain a tight lower bound for the optimal value of the general QOP. Section 4 presents a bisection method together with the proximal alternating direction multiplier method [14] and the accelerated proximal gradient method [4] for solving the Lagrangian-DNN relaxation of the general QOP. Numerical results on the binary integer quadratic problem, the maximum stable set problem, the quadratic multiple knapsack problem, and the quadratic assignment problem are presented in Section 5. Finally, we conclude in Section 6.

2 Preliminaries

2.1 Notation and symbols

We use the following notation and symbols throughout the paper.

$$\begin{aligned}
\mathbb{R}^n &= \text{the space of } n\text{-dimensional column vectors,} \\
\mathbb{R}_+^n &= \text{the nonnegative orthant of } \mathbb{R}^n, \\
\mathbb{S}^n &= \text{the space of } n \times n \text{ symmetric matrices,} \\
\mathbb{S}_+^n &= \text{the cone of } n \times n \text{ symmetric positive semidefinite matrices,} \\
\mathbb{C} &= \{ \mathbf{A} \in \mathbb{S}^n : \mathbf{x}^T \mathbf{A} \mathbf{x} \geq 0 \text{ for all } \mathbf{x} \in \mathbb{R}_+^n \} \text{ (the copositive cone),} \\
\mathbf{\Gamma} &= \{ \mathbf{x} \mathbf{x}^T : \mathbf{x} \in \mathbb{R}_+^n \}, \\
\mathbb{C}^* &= \text{the convex hull of } \mathbf{\Gamma} \text{ (the completely positive cone, the dual of } \mathbb{C} \text{),} \\
\mathbb{N} &= \text{the space of } n \times n \text{ symmetric matrices with nonnegative elements,} \\
\mathbf{Y} \bullet \mathbf{Z} &= \text{trace of } \mathbf{Y} \mathbf{Z} \text{ for every } \mathbf{Y}, \mathbf{Z} \in \mathbb{S}^n \text{ (the inner product).}
\end{aligned}$$

The following relations for $\mathbf{\Gamma}$, \mathbb{S}_+^n , \mathbb{C} , \mathbb{C}^* and \mathbb{N} are well-known:

$$\begin{aligned}
\mathbf{\Gamma} &\subset \mathbb{C}^* \subset \mathbb{S}_+^n \cap \mathbb{N} \subset \mathbb{S}_+^n \subset \mathbb{S}_+^n + \mathbb{N} \subset \mathbb{C}, \\
\mathbf{\Gamma} &= \{ \mathbf{X} \in \mathbb{S}_+^n \cap \mathbb{N} : \text{rank}(\mathbf{X}) = 1 \}, \\
\mathbb{S}_+^n \cap \mathbb{N} &= (\mathbb{S}_+^n + \mathbb{N})^* \text{ (the dual of } \mathbb{S}_+^n + \mathbb{N} \text{).}
\end{aligned}$$

We call $\mathbb{S}_+^n \cap \mathbb{N}$ *the doubly nonnegative cone*.

For $\mathbf{x} \in \mathbb{R}^n$, \mathbf{x}^T denotes the transpose of \mathbf{x} . We use the notation $(\mathbf{t}, \mathbf{u}) \in \mathbb{R}^{\ell+m}$ for the $(\ell + m)$ -dimensional column vector consisting of $\mathbf{t} \in \mathbb{R}^\ell$ and $\mathbf{u} \in \mathbb{R}^m$. In the subsequent discussions, the quadratic form $\mathbf{x}^T \mathbf{Q} \mathbf{x}$ associated with a matrix $\mathbf{Q} \in \mathbb{S}^n$ is represented as $\mathbf{Q} \bullet \mathbf{x} \mathbf{x}^T$ to suggest that $\mathbf{Q} \bullet \mathbf{x} \mathbf{x}^T$ with $\mathbf{x} \in \mathbb{R}_+^n$ is relaxed to $\mathbf{Q} \bullet \mathbf{X}$ with $\mathbf{X} \in \mathbb{C}^*$, $\mathbf{X} \in \mathbb{S}_+^n \cap \mathbb{N}$ or $\mathbf{X} \in \mathbb{S}_+^n$.

2.2 A quadratic optimization model

We introduce a QOP of the form

$$\zeta := \inf \left\{ \mathbf{Q}_0 \bullet \mathbf{x}\mathbf{x}^T \mid \begin{array}{l} \mathbf{x} \in \mathbb{R}_+^n, \mathbf{H}_0 \bullet \mathbf{x}\mathbf{x}^T = 1, \\ \mathbf{Q}_p \bullet \mathbf{x}\mathbf{x}^T = 0 \ (p = 1, 2, \dots, q) \end{array} \right\}, \quad (2)$$

where $\mathbf{H}_0 \in \mathbb{S}^n$ and $\mathbf{Q}_p \in \mathbb{S}^n$ ($p = 0, 1, \dots, q$), to describe a class of conic relaxations and their further Lagrangian relaxations in Section 3. This form of QOP was introduced in [1] to establish an equivalence to its CPP relaxation under a set of conditions. If we are concerned with only the CPP relaxation among the conic relaxations, basic theoretical results presented in Section 3 can be obtained from [1] and [2]. Our main emphasis here is on extending their framework to a larger class of conic and Lagrangian-conic relaxations. In particular, the class includes a Lagrangian-DNN relaxation of QOP (2), which is shown to work very effectively and efficiently for large-scale QOPs in Section 5 with the first order methods described in Section 4.

We assume the following three conditions throughout the paper. These conditions are stronger than the ones assumed in [1] and [2], because the framework includes not only the CPP relaxation but also the DNN relaxation.

Condition (a) The feasible region

$$\{\mathbf{x} \in \mathbb{R}_+^n : \mathbf{H}_0 \bullet \mathbf{x}\mathbf{x}^T = 1, \mathbf{Q}_p \bullet \mathbf{x}\mathbf{x}^T = 0 \ (p = 1, 2, \dots, q)\} \quad (3)$$

is nonempty.

Condition (b) $\mathbf{O} \neq \mathbf{H}_0 \in \mathbb{S}_+^n + \mathbb{N}$ and $\mathbf{O} \neq \mathbf{Q}_p \in \mathbb{S}_+^n + \mathbb{N}$ ($p = 1, 2, \dots, q$).

Condition (c) $\mathbf{D} = \mathbf{O}$ if $\mathbf{D} \in \mathbb{S}_+^n \cap \mathbb{N}$, $\mathbf{H}_0 \bullet \mathbf{D} = 0$ and $\mathbf{Q}_p \bullet \mathbf{D} = 0$ ($p = 1, 2, \dots, q$).

Notice that if $\mathbf{H}_0 = \mathbf{O}$ then QOP (2) is infeasible, and if $\mathbf{Q}_p = \mathbf{O}$ for some p then the redundant constraint $\mathbf{Q}_p \bullet \mathbf{x}\mathbf{x}^T = 0$ can be eliminated. Thus, $\mathbf{H}_0 \neq \mathbf{O}$ and $\mathbf{Q}_p \neq \mathbf{O}$ ($p = 1, 2, \dots, q$) in Condition (b) can be assumed without loss of generality. We note that Condition (c) together with Condition (a) require that the feasible region is nonempty and bounded. For the proof of this fact, see (i) of Lemma 3.1 and its proof. Hence “inf” in (2) can be replaced by “min”. We let \mathbf{x}^* be an optimal solution of QOP (2) with the finite optimal value ζ .

We can easily transform QOP (1) with linear and complementarity constraints to a

QOP of the form (2) satisfying Conditions (a), (b) and (c) as follows. Define

$$\begin{aligned}
\mathbf{x} &= (x_1, \mathbf{u}) \in \mathbb{R}^n \text{ with } n = 1 + m, \\
\mathbf{Q}_0 &= \begin{pmatrix} 0 & \mathbf{c}^T \\ \mathbf{c} & \mathbf{Q} \end{pmatrix} \in \mathbb{S}^n, \quad \mathbf{H}_0 = \begin{pmatrix} 1 & \mathbf{0}^T \\ \mathbf{0} & \mathbf{O} \end{pmatrix} \in \mathbb{S}^n, \\
\mathbf{Q}_{01} &= \begin{pmatrix} \mathbf{b}^T \mathbf{b} & \mathbf{b}^T \mathbf{A} \\ \mathbf{A}^T \mathbf{b} & \mathbf{A}^T \mathbf{A} \end{pmatrix} \in \mathbb{S}^n, \\
\mathbf{C}_{ij} &= \text{the } m \times m \text{ matrix with } (i, j)\text{th component } 1/2 \\
&\quad \text{and } 0 \text{ elsewhere } ((i, j) \in \mathcal{E}), \\
\mathbf{Q}_{ij} &= \begin{pmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & \mathbf{C}_{ij} + \mathbf{C}_{ij}^T \end{pmatrix} \in \mathbb{S}^n \quad ((i, j) \in \mathcal{E}).
\end{aligned}$$

Then QOP (1) is equivalent to

$$\text{minimize } \left\{ \mathbf{Q}_0 \bullet \mathbf{x}\mathbf{x}^T \mid \begin{array}{l} \mathbf{x} \in \mathbb{R}_+^n, \mathbf{H}_0 \bullet \mathbf{x}\mathbf{x}^T = 1, \\ \mathbf{Q}_{01} \bullet \mathbf{x}\mathbf{x}^T = 0, \mathbf{Q}_{ij} \bullet \mathbf{x}\mathbf{x}^T = 0 \quad ((i, j) \in \mathcal{E}) \end{array} \right\} \quad (4)$$

in the sense that $\mathbf{u} \in \mathbb{R}^m$ is a feasible solution of (1) with the objective value $\mathbf{u}^T \mathbf{Q}\mathbf{u} + 2\mathbf{c}^T \mathbf{u}$ if and only if $\mathbf{x} = (1, \mathbf{u}) \in \mathbb{R}^n$ is a feasible solution of (4) with the same objective value $\mathbf{Q}_0 \bullet \mathbf{x}\mathbf{x}^T$.

Lemma 2.1. *Assume that the feasible region of QOP (1) is nonempty and that the polyhedral set $\{\mathbf{u} \in \mathbb{R}_+^m : \mathbf{A}\mathbf{u} + \mathbf{b} = \mathbf{0}\}$ is bounded. Then QOP (4) induced from QOP (1) satisfies Conditions (a), (b) and (c). Here we assume that the subscripts of the matrices \mathbf{Q}_{01} and \mathbf{Q}_{ij} $((i, j) \in \mathcal{E})$ have been renumbered to $1, 2, \dots, q$ for some q .*

Proof. We only prove Condition (c) because Conditions (a) and (b) are obvious. Assume that $\mathbf{H}_0 \bullet \mathbf{D} = 0$, $\mathbf{Q}_{01} \bullet \mathbf{D} = 0$ and $\mathbf{Q}_{ij} \bullet \mathbf{D} = 0$ $(i, j) \in \mathcal{E}$ for some $\mathbf{D} \in \mathbb{S}_+^{1+m} \cap \mathbb{N}$. Then, we see that

$$0 = \mathbf{H}_0 \bullet \mathbf{D} = D_{11}, \quad \text{and } 0 = \mathbf{Q}_{01} \bullet \mathbf{D} = \begin{pmatrix} \mathbf{b}^T \mathbf{b} & \mathbf{b}^T \mathbf{A} \\ \mathbf{A}^T \mathbf{b} & \mathbf{A}^T \mathbf{A} \end{pmatrix} \bullet \mathbf{D}. \quad (5)$$

Now we write $\mathbf{D} \in \mathbb{S}_+^{1+m} \cap \mathbb{N}$ as $\begin{pmatrix} D_{11} & \mathbf{D}_{12} \\ \mathbf{D}_{12}^T & \mathbf{D}_{22} \end{pmatrix}$. From $0 = D_{11}$ and $\mathbf{D} \in \mathbb{S}_+^{1+m}$, we get $\mathbf{D}_{12} = \mathbf{0}$. As a result, the last relation in (5) implies that $\mathbf{A}^T \mathbf{A} \bullet \mathbf{D}_{22} = \mathbf{0}$. Since $\mathbf{A}^T \mathbf{A} \in \mathbb{S}_+^m$ and $\mathbf{D}_{22} \in \mathbb{S}_+^m$, we obtain $\mathbf{A}^T \mathbf{A} \mathbf{D}_{22} = \mathbf{0}$ and $\mathbf{A} \mathbf{D}_{22} = \mathbf{0}$. On the other hand, by the assumption of the lemma, there does not exist nonzero $\mathbf{d} \in \mathbb{R}^m$ such that $\mathbf{d} \geq \mathbf{0}$ and $-\mathbf{A}\mathbf{d} = \mathbf{0}$. By applying Tucker's theorem of the alternative [30], we can take a $\mathbf{y} \in \mathbb{R}^m$ such that $\mathbf{A}^T \mathbf{y} > \mathbf{0}$. Multiplying \mathbf{y}^T to the identity $\mathbf{A} \mathbf{D}_{22} = \mathbf{0}$, we obtain that $(\mathbf{y}^T \mathbf{A}) \mathbf{D}_{22} = \mathbf{0}^T$, $(\mathbf{y}^T \mathbf{A}) > \mathbf{0}^T$ and $\mathbf{D}_{22} \in \mathbb{N}$, which implies $\mathbf{D}_{22} = \mathbf{0}$. Thus we have shown that $\mathbf{D} = \mathbf{0}$. \square

Condition (b) implies that the inequalities $\mathbf{Q}_p \bullet \mathbf{x}\mathbf{x}^T \geq 0$ ($p = 1, 2, \dots, q$) hold for any $\mathbf{x} \in \mathbb{R}_+^n$. Hence, the set of equalities $\mathbf{Q}_p \bullet \mathbf{x}\mathbf{x}^T = 0$ ($p = 1, 2, \dots, q$) in QOP (2) can be combined into a single equality $\mathbf{H}_1 \bullet \mathbf{x}\mathbf{x}^T = 0$, where $\mathbf{H}_1 = \sum_{p=1}^q \mathbf{Q}_p$. Consequently, we obtain a simplified QOP:

$$\zeta := \min \left\{ \mathbf{Q}_0 \bullet \mathbf{x}\mathbf{x}^T \mid \mathbf{x} \in \mathbb{R}_+^n, \mathbf{H}_0 \bullet \mathbf{x}\mathbf{x}^T = 1, \mathbf{H}_1 \bullet \mathbf{x}\mathbf{x}^T = 0 \right\}, \quad (6)$$

which is equivalent to QOP (2). Specifically, (2) and (6) share a common feasible region (3) and a common optimal solution \mathbf{x}^* with the optimal value ζ . We note that QOP (4) (hence (1)) is reduced to QOP (6) if we define $\mathbf{H}_1 = \mathbf{Q}_{01} + \sum_{(i,j) \in \mathcal{E}} \mathbf{Q}_{ij}$.

3 Main results

We present a class of conic relaxations of QOP (6), which includes the CPP and DNN relaxations of QOP (6), in Section 3.1, and their further Lagrangian relaxations, called Lagrangian-conic relaxations, in Section 3.2. From the theoretical results shown in Lemmas 3.1, 3.2 and 3.3, we can conclude that the Lagrangian-DNN relaxation of QOP (6) is almost as effective as the DNN relaxation applied to the original QOP (2), and that solving the Lagrangian-DNN relaxation is expected to be more efficient and stable than solving the direct DNN relaxation of (2); see also Remark 3.1.

3.1 A class of conic relaxations

We rewrite QOP (2) as

$$\zeta := \min \left\{ \mathbf{Q}_0 \bullet \mathbf{X} \mid \begin{array}{l} \mathbf{X} \in \Gamma, \mathbf{H}_0 \bullet \mathbf{X} = 1, \\ \mathbf{Q}_p \bullet \mathbf{X} = 0 \ (p = 1, 2, \dots, q) \end{array} \right\}$$

and QOP (6) as

$$\zeta := \min \{ \mathbf{Q}_0 \bullet \mathbf{X} \mid \mathbf{H}_0 \bullet \mathbf{X} = 1, \mathbf{H}_1 \bullet \mathbf{X} = 0, \mathbf{X} \in \Gamma \}.$$

Recall that $\Gamma = \{ \mathbf{x}\mathbf{x}^T : \mathbf{x} \in \mathbb{R}_+^n \}$. If Γ is replaced by a closed convex cone \mathbb{K} in \mathbb{S}^n satisfying $\Gamma \subset \mathbb{K}$, then the following convex relaxations of QOPs (2) and (6) are obtained:

$$\zeta(\mathbb{K}) := \inf \left\{ \mathbf{Q}_0 \bullet \mathbf{X} \mid \begin{array}{l} \mathbf{X} \in \mathbb{K}, \mathbf{H}_0 \bullet \mathbf{X} = 1, \\ \mathbf{Q}_p \bullet \mathbf{X} = 0 \ (p = 1, 2, \dots, q) \end{array} \right\} \quad (7)$$

$$\eta(\mathbb{K}) := \inf \{ \mathbf{Q}_0 \bullet \mathbf{X} \mid \mathbf{H}_0 \bullet \mathbf{X} = 1, \mathbf{H}_1 \bullet \mathbf{X} = 0, \mathbf{X} \in \mathbb{K} \}. \quad (8)$$

Note that dual problem of (8) is given by

$$\eta^d(\mathbb{K}) := \sup \left\{ y_0 \mid \mathbf{Z} + y_0 \mathbf{H}_0 + y_1 \mathbf{H}_1 = \mathbf{Q}_0, \mathbf{Z} \in \mathbb{K}^*, \mathbf{y} = (y_0, y_1) \in \mathbb{R}^2 \right\}. \quad (9)$$

When \mathbb{K} is chosen to be \mathbb{C}^* , $\mathbb{S}_+^n \cap \mathbb{N}$ or \mathbb{S}_+^n , the problem (7) (or the problem (8)) is known as a CPP relaxation, a DNN relaxation and an SDP relaxation of QOP (2) (or QOP (6)), respectively. As the CPP relaxation attains the exact optimal value ζ of QOP (2) (or QOP (6)) (see Theorem 3.5 of [1]), it is theoretically the most important among the three relaxations. It is, however, numerically intractable while the DNN and SDP relaxations are numerically implementable. From $\mathbf{\Gamma} \subset \mathbb{C}^* \subset \mathbb{S}_+^n \cap \mathbb{N} \subset \mathbb{S}_+^n$ and

$$\zeta(\mathbb{S}_+^n) \leq \zeta(\mathbb{S}_+^n \cap \mathbb{N}) \leq \zeta(\mathbb{C}^*) = \zeta(\mathbf{\Gamma}) = \zeta,$$

the DNN relaxation of QOP (2) provides a lower bound for the minimum value ζ of (2) at least as effectively as the SDP relaxation. Furthermore, under Conditions (a), (b) and (c), the Lagrangian-DNN relaxation of (6) (for which (2) is equivalent to) satisfies additional properties which are conducive for numerically solving the problem, especially with first-order methods [4, 32].

We present three lemmas to show those properties in the remainder of Section 3 for the general case \mathbb{K} where $\mathbf{\Gamma} \subset \mathbb{K} \subset \mathbb{S}_+^n \cap \mathbb{N}$. They are significant in their own right, although a numerical method is proposed only for $\mathbb{K} = \mathbb{S}_+^n \cap \mathbb{N}$ in Section 4.

Lemma 3.1. *Suppose that \mathbb{K} is a closed (not necessarily convex) cone in \mathbb{S}^n satisfying $\mathbf{\Gamma} \subset \mathbb{K} \subset \mathbb{S}_+^n \cap \mathbb{N}$.*

- (i) *The feasible region of the problem (7) with \mathbb{K} is nonempty and bounded; hence $\zeta(\mathbb{K}) > -\infty$.*
- (ii) *$\eta(\mathbb{K}) = \zeta(\mathbb{K}) \leq \zeta$.*

Proof. By Condition (a), we know that $\mathbf{x}^*(\mathbf{x}^*)^T$ is a feasible solution of the problem (7) with \mathbb{K} . For assertion (i), it suffices to show that the feasible region of the problem (7) with \mathbb{K} is bounded. Assume on the contrary that there exists a sequence $\{\mathbf{X}^k \in \mathbb{K}\}$ with $\lim_{k \rightarrow \infty} \|\mathbf{X}^k\| = \infty$ such that

$$\mathbf{H}_0 \bullet \mathbf{X}^k = 1 \text{ and } \mathbf{Q}_p \bullet \mathbf{X}^k = 0 \text{ (} p = 1, 2, \dots, q \text{)}.$$

We may assume without loss of generality that $\mathbf{X}^k / \|\mathbf{X}^k\|$ converges to some $\mathbf{D} \in \mathbb{K} \subset \mathbb{S}_+^n \cap \mathbb{N}$. Dividing the identities above by $\|\mathbf{X}^k\|$ and taking their limit as $k \rightarrow \infty$, we have that

$$\mathbf{O} \neq \mathbf{D} \in \mathbb{K} \subset \mathbb{S}_+^n \cap \mathbb{N}, \mathbf{H}_0 \bullet \mathbf{D} = 0 \text{ and } \mathbf{Q}_p \bullet \mathbf{D} = 0 \text{ (} p = 1, 2, \dots, q \text{)}.$$

This contradicts the given Condition (c), and we have shown assertion (i).

By the assumption, $\mathbf{\Gamma} \subset \mathbb{K}$. Hence $\zeta(\mathbb{K}) \leq \zeta(\mathbf{\Gamma}) = \zeta$. Since $\mathbf{H}_1 = \sum_{p=1}^q \mathbf{Q}_p$, if $\mathbf{X} \in \mathbb{K}$ is a feasible solution of (7) with the objective value $\mathbf{Q}_0 \bullet \mathbf{X}$, then it is a feasible solution of (8) with the same objective value. Thus, $\eta(\mathbb{K}) \leq \zeta(\mathbb{K})$ follows. To prove the converse inequality, suppose that $\mathbf{X} \in \mathbb{K}$ is a feasible solution of (8) with $\mathbb{K} \subset \mathbb{S}_+^n \cap \mathbb{N}$. By Condition

(b), we know $\mathbf{Q}_p \in \mathbb{S}_+^n + \mathbb{N}$ ($p = 1, 2, \dots, q$). Hence $\mathbf{Q}_p \bullet \mathbf{X} \geq 0$ ($p = 1, 2, \dots, q$), which together with $0 = \mathbf{H}_1 \bullet \mathbf{X} = \sum_{p=1}^q (\mathbf{Q}_p \bullet \mathbf{X})$ imply that $\mathbf{Q}_p \bullet \mathbf{X} = 0$ ($p = 1, 2, \dots, q$). Therefore, \mathbf{X} is a feasible solution of (7) with the objective value $\mathbf{Q} \bullet \mathbf{X}$, and the inequality $\eta(\mathbb{K}) \geq \zeta(\mathbb{K})$ follows. \square

Observe that if the relaxation technique discussed above is directly applied to QOP (1), then we have the following problem:

$$\text{minimize } \left\{ \mathbf{Q} \bullet \mathbf{U} + 2\mathbf{c}^T \mathbf{u} \mid \begin{array}{l} \mathbf{X} = \begin{pmatrix} 1 & \mathbf{u}^T \\ \mathbf{u} & \mathbf{U} \end{pmatrix} \in \mathbb{K}, \mathbf{A}\mathbf{u} + \mathbf{b} = \mathbf{0}, \\ \mathbf{Q}_{ij} \bullet \mathbf{X} = 0 \ ((i, j) \in \mathcal{E}) \end{array} \right\}. \quad (10)$$

For $\mathbb{K} = \mathbf{\Gamma}$, the problems (7) induced from (4) and (10) induced from (1) are equivalent, and both problems represent QOP (1). If we choose a closed convex cone \mathbb{K} with $\mathbf{\Gamma} \subset \mathbb{K}$, both problems serve as convex relaxations of QOP (1), but they are not equivalent in general. The essential difference lies in $\mathbf{A}\mathbf{u} + \mathbf{b} = \mathbf{0}$ and $\mathbf{Q}_{01} \bullet \mathbf{X} = 0$. Suppose that $\mathbf{X} = \begin{pmatrix} 1 & \mathbf{u}^T \\ \mathbf{u} & \mathbf{U} \end{pmatrix} \in \mathbb{K}$ is a feasible solution of the problem (7) with $\mathbf{\Gamma} \subset \mathbb{K} \subset \mathbb{S}_+^n$. Then it satisfies

$$0 = \mathbf{Q}_{01} \bullet \begin{pmatrix} 1 & \mathbf{u}^T \\ \mathbf{u} & \mathbf{U} \end{pmatrix} = \begin{pmatrix} \mathbf{b}^T \mathbf{b} & \mathbf{b}^T \mathbf{A} \\ \mathbf{A}^T \mathbf{b} & \mathbf{A}^T \mathbf{A} \end{pmatrix} \bullet \begin{pmatrix} 1 & \mathbf{u}^T \\ \mathbf{u} & \mathbf{U} \end{pmatrix}.$$

Since $\mathbf{Q}_{01} \in \mathbb{S}_+^n$ and $\mathbf{X} \in \mathbb{S}_+^n$, we see that $\mathbf{Q}_{01} \mathbf{X} = \mathbf{O}$. It follows that

$$\mathbf{A}\mathbf{u} + \mathbf{b} = \mathbf{0} \quad \text{and} \quad \mathbf{b}\mathbf{u}^T + \mathbf{A}\mathbf{U} = \mathbf{O}.$$

From the first equality above, we know that \mathbf{X} is a feasible solution of the problem (10); hence the problem (7) (hence (8)) provides a convex relaxation at least as good as the problem (10). Furthermore, the second equality, which is not involved in (10) unless $\text{rank}(\mathbf{X}) = 1$, often contributes to strengthening the relaxation.

3.2 A class of Lagrangian-conic relaxations

For each closed cone \mathbb{K} (not necessarily convex), we consider a Lagrangian relaxation of the problem (8) and its dual.

$$\eta^p(\lambda, \mathbb{K}) := \inf \{ \mathbf{Q}_0 \bullet \mathbf{X} + \lambda \mathbf{H}_1 \bullet \mathbf{X} \mid \mathbf{H}_0 \bullet \mathbf{X} = 1, \mathbf{X} \in \mathbb{K} \}, \quad (11)$$

$$\eta^d(\lambda, \mathbb{K}) := \sup \{ y_0 \mid \mathbf{Q}_0 + \lambda \mathbf{H}_1 - y_0 \mathbf{H}_0 \in \mathbb{K}^* \}, \quad (12)$$

where $\lambda \in \mathbb{R}$ denotes a Lagrangian parameter. We call either of (11) and (12) a Lagrangian-conic relaxation of QOP (6), and particularly a Lagrangian-DNN relaxation when $\mathbb{K} = \mathbb{S}^n \cap \mathbb{N}$. (We use these names for simplicity, although (12) is precisely the dual of

Lagrangian-conic or Lagrangian-DNN relaxation.) It is easily verified that the weak duality relation $\eta^d(\lambda, \mathbb{K}) \leq \eta^p(\lambda, \mathbb{K}) \leq \eta(\mathbb{K})$ hold for every $\lambda \in \mathbb{R}$. Note that by Condition (b), the problem (11) is strictly feasible when \mathbb{K} includes the completely positive cone \mathbb{C}^* with nonempty interior w.r.t. \mathbb{S}^n (see, for example, [12]).

Suppose that $\Gamma \subset \mathbb{K} \subset \mathbb{S}_+^n \cap \mathbb{N}$. Then we see by Condition (b) that $\mathbf{H}_1 \in \mathbb{S}_+^n + \mathbb{N}$. Hence $\mathbf{H}_1 \bullet \mathbf{X} \geq 0$ for every $\mathbf{X} \in \mathbb{K}$. Thus, the second term $\lambda \mathbf{H}_1 \bullet \mathbf{X}$ of the objective function of (11) serves as a penalty function for the equality constraint $\mathbf{H}_1 \bullet \mathbf{X} = 0$ such that for each $\mathbf{X} \in \mathbb{K}$ and $\lambda \geq 0$,

$$\lambda \mathbf{H}_1 \bullet \mathbf{X} \geq 0 \quad \text{and} \quad \lambda \mathbf{H}_1 \bullet \mathbf{X} \rightarrow \infty \quad \text{as} \quad \lambda \rightarrow \infty \quad \text{if and only if} \quad \mathbf{H}_1 \bullet \mathbf{X} \neq 0. \quad (13)$$

By Lemma 3.1, we also know that $-\infty < \eta(\mathbb{K}) = \zeta(\mathbb{K}) \leq \zeta$. Using these relations, we establish the following result.

Lemma 3.2. *Suppose that \mathbb{K} is a closed cone (not necessarily convex) in \mathbb{S}^n satisfying $\Gamma \subset \mathbb{K} \subset \mathbb{S}_+^n \cap \mathbb{N}$. Then the following statements hold.*

(i) $\eta^p(\lambda_1, \mathbb{K}) \leq \eta^p(\lambda_2, \mathbb{K}) \leq \eta(\mathbb{K})$ if $0 < \lambda_1 < \lambda_2$.

(ii) $\eta^d(\lambda_1, \mathbb{K}) \leq \eta^d(\lambda_2, \mathbb{K}) \leq \eta(\mathbb{K})$ if $0 < \lambda_1 < \lambda_2$.

(iii) $\eta^p(\lambda, \mathbb{K})$ converges to $\eta(\mathbb{K})$ as $\lambda \rightarrow \infty$.

(iv) Moreover, if \mathbb{K} is convex, then $\eta^d(\lambda, \mathbb{K}) = \eta^p(\lambda, \mathbb{K})$ for every large $\lambda > 0$.

Proof. Assertion (i) follows from the first relation of (13) and the weak duality of Lagrangian relaxation. Assertion (ii) follows from the fact that $\mathbf{H}_1 \in \mathbb{S}_+^n + \mathbb{N} \subset \mathbb{K}^* \subset \Gamma^* = \mathbb{C}$. To prove (iii), define the level set with the objective value $\eta(\mathbb{K}) > -\infty$ by

$$L(\lambda, \mathbb{K}) = \{ \mathbf{X} \in \mathbb{K} : \mathbf{H}_0 \bullet \mathbf{X} = 1, \mathbf{Q}_0 \bullet \mathbf{X} + \lambda \mathbf{H}_1 \bullet \mathbf{X} \leq \eta(\mathbb{K}) \}$$

for the problem (11) with each $\lambda \geq 0$. Then $L(\lambda, \mathbb{K})$ contains an optimal solution $\widetilde{\mathbf{X}}$ of (8) for every $\lambda \geq 0$, and $L(\lambda_1, \mathbb{K}) \supset L(\lambda_2, \mathbb{K})$ if $0 < \lambda_1 < \lambda_2$. We will show that $L(\lambda, \mathbb{K})$ is bounded for a sufficiently large $\lambda > 0$. Assume on the contrary that there exists a sequence $\{(\lambda^k, \mathbf{X}^k) \in \mathbb{R}_+ \times \mathbb{K}\}$ such that $\mathbf{X}^k \in L(\lambda^k, \mathbb{K})$, $0 < \lambda^k \rightarrow \infty$ and $0 < \|\mathbf{X}^k\| \rightarrow \infty$ as $k \rightarrow \infty$. Then, we have

$$\begin{aligned} \frac{\mathbf{X}^k}{\|\mathbf{X}^k\|} &\in \mathbb{K}, \quad \mathbf{H}_1 \bullet \frac{\mathbf{X}^k}{\|\mathbf{X}^k\|} \geq 0 \quad (\text{by } \mathbf{H}_1 \in \mathbb{S}_+^n + \mathbb{N}), \\ \mathbf{H}_0 \bullet \frac{\mathbf{X}^k}{\|\mathbf{X}^k\|} &= 0 \quad \text{and} \quad \mathbf{Q}_0 \bullet \frac{\mathbf{X}^k}{\lambda^k \|\mathbf{X}^k\|} + \mathbf{H}_1 \bullet \frac{\mathbf{X}^k}{\|\mathbf{X}^k\|} \leq \frac{\zeta}{\lambda^k \|\mathbf{X}^k\|} \end{aligned}$$

We may assume without loss of generality that $\mathbf{X}^k / \|\mathbf{X}^k\|$ converges to a nonzero $\mathbf{D} \in \mathbb{K}$. By taking the limit as $k \rightarrow \infty$, we obtain that

$$\mathbf{0} \neq \mathbf{D} \in \mathbb{K} \subset \mathbb{S}_+^n \cap \mathbb{N}, \quad \mathbf{H}_0 \bullet \mathbf{D} = 0, \quad \mathbf{H}_1 \bullet \mathbf{D} = 0.$$

This contradicts the given Condition (c). Therefore, we have shown that $L(\bar{\lambda}, \mathbb{K})$ is bounded for some sufficiently large $\bar{\lambda} > 0$ and $\widetilde{\mathbf{X}} \in L(\lambda, \mathbb{K}) \subset L(\bar{\lambda}, \mathbb{K})$ for every $\lambda \geq \bar{\lambda}$.

Let $\{\lambda^k \geq \bar{\lambda}\}$ be a sequence that diverges to ∞ . Since the nonempty level set $L(\lambda^k, \mathbb{K})$ is contained in a bounded set $L(\bar{\lambda}, \mathbb{K})$, the problem (11) with each $\lambda = \lambda^k$ has an optimal solution \mathbf{X}^k with the objective value $\eta^p(\lambda^k, \mathbb{K}) = \mathbf{Q}_0 \bullet \mathbf{X}^k + \lambda^k \mathbf{H}_1 \bullet \mathbf{X}^k$ in the level set $L(\lambda^k, \mathbb{K})$. We may assume without loss of generality that \mathbf{X}^k converges to some $\bar{\mathbf{X}} \in L(\bar{\lambda}, \mathbb{K})$. It follows that

$$\begin{aligned} \mathbf{H}_0 \bullet \mathbf{X}^k &= 1, \quad \frac{\mathbf{Q}_0 \bullet \mathbf{X}^k}{\lambda^k} + \mathbf{H}_1 \bullet \mathbf{X}^k \leq \frac{\eta(\mathbb{K})}{\lambda^k}, \\ \mathbf{Q}_0 \bullet \mathbf{X}^k &\leq \eta(\mathbb{K}), \quad \mathbf{H}_1 \bullet \mathbf{X}^k \geq 0 \quad (\text{both by Condition (b)}). \end{aligned}$$

By taking the limit as $k \rightarrow \infty$, we obtain that

$$\bar{\mathbf{X}} \in \mathbb{K}, \quad \mathbf{H}_0 \bullet \bar{\mathbf{X}} = 1, \quad \mathbf{H}_1 \bullet \bar{\mathbf{X}} = 0, \quad \mathbf{Q}_0 \bullet \bar{\mathbf{X}} \leq \eta(\mathbb{K}).$$

This implies that $\bar{\mathbf{X}}$ is an optimal solution of the problem (8). Hence, $\mathbf{Q}_0 \bullet \mathbf{X}^k$ converges to $\eta(\mathbb{K})$ as $k \rightarrow \infty$. We also see from

$$\mathbf{Q}_0 \bullet \mathbf{X}^k \leq \eta^p(\lambda^k, \mathbb{K}) = \mathbf{Q}_0 \bullet \mathbf{X}^k + \lambda^k \mathbf{H}_1 \bullet \mathbf{X}^k \leq \eta(\mathbb{K})$$

that $\eta^p(\lambda^k, \mathbb{K})$ converges to $\eta(\mathbb{K})$. Thus, we have shown assertion (iii).

Finally, we prove assertion (iv). We first see that the problem (11) has an interior feasible solution by Condition (b). Indeed, if $\widetilde{\mathbf{X}}$ is an interior point of \mathbb{C}^* , which also lies in the interior of any closed convex cone \mathbb{K} satisfying $\mathbf{\Gamma} \subset \mathbb{K} \subset \mathbb{S}_+^n \cap \mathbb{N}$, then $\mathbf{H}_0 \bullet \widetilde{\mathbf{X}} > 0$. Hence $\widetilde{\mathbf{X}} / (\mathbf{H}_0 \bullet \widetilde{\mathbf{X}})$ serves as an interior feasible solution of the problems (11) with \mathbb{K} . On the other hand, we have observed in the proof of assertion (iii) above that the problem (11) has optimal solutions if $\lambda > \bar{\lambda}$. Therefore, by the duality theorem for linear optimization problems over closed convex cones (see, for example, Theorem 4.2.1 of [22]), we obtain that $\eta^d(\lambda, \mathbb{K}) = \eta^p(\lambda, \mathbb{K})$ for every $\lambda \geq \bar{\lambda}$. \square

Lemma 3.3. *Suppose that \mathbb{K} is a closed convex cone in \mathbb{S}^n satisfying $\mathbf{\Gamma} \subset \mathbb{K} \subset \mathbb{S}_+^n \cap \mathbb{N}$.*

- (i) *Suppose $\eta^p(\lambda, \mathbb{K}) < \eta(\mathbb{K})$ for all λ , i.e., $\eta(\mathbb{K})$ is not attained by $\eta^p(\lambda^*, \mathbb{K})$ for any finite λ^* . Then the optimal value of the problem (9) is not attained at any feasible solution of the problem.*
- (ii) *Suppose that $q \geq 1$. Then the feasible set of (8), which is nonempty by Lemma 3.1, has no feasible point in the interior of \mathbb{K} .*

Proof. (i) We proof the result by contradiction. Suppose (9) attained the optimal value at a $\mathbf{y}^* = (y_0^*, y_1^*) \in \mathbb{R}^n$ with $y_0^* = \eta^d(\mathbb{K})$. By definition, it is clear that $\eta^d(-y_1^*, \mathbb{K}) \leq \eta^d(\mathbb{K})$.

On the other hand, since y_0^* is feasible for (12) with parameter equal to $-y_1^*$, we have $\eta^d(-y_1^*, \mathbb{K}) \geq y_0^*$. Thus $y_0^* = \eta^d(\mathbb{K}) = \eta^d(-y_1^*, \mathbb{K})$. Now for $\lambda \geq -y_1^*$, we have

$$\eta^d(\mathbb{K}) = \eta^d(-y_1^*, \mathbb{K}) \leq \eta^d(\lambda, \mathbb{K}) \leq \eta^d(\mathbb{K}).$$

Hence $\eta^d(\mathbb{K}) = \eta^d(-y_1^*, \mathbb{K}) = \eta^d(\lambda, \mathbb{K}) = \eta^p(\lambda, \mathbb{K})$ for every sufficiently large $\lambda \geq -y_1^*$, where the last equality follows from assertion (iv) in Lemma 3.2. By letting $\lambda \uparrow \infty$, we get by using assertion (iii) of Lemma 3.2 that $\eta^d(-y_1^*, \mathbb{K}) = \eta(\mathbb{K})$. Since $\eta^d(-y_1^*, \mathbb{K}) \leq \eta^p(-y_1^*, \mathbb{K}) \leq \eta(\mathbb{K})$, we deduce that $\eta^p(-y_1^*, \mathbb{K}) = \eta(\mathbb{K})$. But this contradicts the condition that $\eta^p(\lambda, \mathbb{K}) < \eta(\mathbb{K})$ for all λ . Hence the optimal value $\eta^d(\mathbb{K})$ is not attained.

(ii) Let \mathbf{X} be an interior point of \mathbb{K} . By Condition (b), we know that $\mathbf{O} \neq \mathbf{Q}_p \in \mathbb{S}_+^n + \mathbb{N}$ ($p = 1, 2, \dots, q$). Since \mathbf{X} lies in the interior of \mathbb{K} , there exists a positive number ϵ such that $\mathbf{X} - \epsilon \mathbf{Q}_p$ ($p = 1, 2, \dots, q$) remain in $\mathbb{K} \subset \mathbb{S}_+^n \cap \mathbb{N}$; hence $\mathbf{Q}_p \bullet (\mathbf{X} - \epsilon \mathbf{Q}_p) \geq 0$ ($p = 1, 2, \dots, q$). It follows that

$$\mathbf{Q}_p \bullet \mathbf{X} > \mathbf{Q}_p \bullet \mathbf{X} - \epsilon \mathbf{Q}_p \bullet \mathbf{Q}_p = \mathbf{Q}_p \bullet (\mathbf{X} - \epsilon \mathbf{Q}_p) \geq 0 \quad (p = 1, 2, \dots, q).$$

Since $q \geq 1$ by the assumption, we obtain that $\mathbf{H}_1 \bullet \mathbf{X} = \sum_{p=1}^q \mathbf{Q}_p \bullet \mathbf{X} > 0$, and any interior point of \mathbb{K} cannot be a feasible solution of (8). \square

Remark 3.1. *The result in Lemma 3.3 has important implications for the numerical solution of the DNN relaxation of QOP (6) obtained as the problem (8) with $\mathbb{K} = \mathbb{S}_+^n \cap \mathbb{N}$ where it has no interior feasible solution. As a slight perturbation of the problem data may render the problem to be infeasible, it is typically expected that a numerical algorithm for solving the problem will encounter numerical difficulties when the iterates approach optimality. In addition, it is generally difficult to compute an accurate approximate optimal solution for such a problem.*

On the other hand, the Lagrangian-DNN relaxation of (6) obtained as the problem (11) with $\mathbb{K} = \mathbb{S}_+^n \cap \mathbb{N}$ is strictly feasible by Condition (b). As a result, the dual problem (12) with $\mathbb{K} = \mathbb{S}_+^n \cap \mathbb{N}$, which we also call the Lagrangian-DNN relaxation, attains its optimal value. These properties are conducive for a numerical algorithm to solve the problems efficiently. Thus in the next section, we will focus on designing an efficient algorithm to solve (12) with $\mathbb{K} = \mathbb{S}_+^n \cap \mathbb{N}$.

4 Algorithms

Here we design an efficient algorithm to solve the problem (12) with $\mathbb{K} = \mathbb{S}_+^n \cap \mathbb{N}$ (the Lagrangian-DNN relaxation of QOP (6)), as mentioned in Remark 3.1. For the subsequent discussion, we let $\mathbb{K}_1 = \mathbb{S}_+^n$ and $\mathbb{K}_2 = \mathbb{N}$, and hence $\mathbb{K}^* = \mathbb{K}_1^* + \mathbb{K}_2^*$. We use $\Pi_{\mathbb{K}}(\cdot)$, $\Pi_{\mathbb{K}^*}(\cdot)$, $\Pi_i(\cdot)$ and $\Pi_i^*(\cdot)$ ($i = 1, 2$) to denote the (metric) projections onto \mathbb{K} , \mathbb{K}^* , \mathbb{K}_i and \mathbb{K}_i^* ($i = 1, 2$), respectively.

Suppose for the moment that we have an efficient algorithm to compute $\Pi_{\mathbb{K}^*}(\mathbf{G})$ for any given $\mathbf{G} \in \mathbb{S}^n$. (Note that to compute $\Pi_{\mathbb{K}^*}(\mathbf{G})$, we will present an accelerated proximal gradient method [4] described as Algorithm C in Section 4.1.) Then we can design a bisection method to solve (12), as we shall describe next.

For a sufficiently large and fixed $\lambda > 0$, define the function

$$g_\lambda(y_0) = \|\mathbf{G}_\lambda(y_0) - \Pi_{\mathbb{K}^*}(\mathbf{G}_\lambda(y_0))\| = \|\Pi_{\mathbb{K}}(-\mathbf{G}_\lambda(y_0))\|$$

where $\mathbf{G}_\lambda(y_0) = \mathbf{Q}_0 + \lambda\mathbf{H}_1 - y_0\mathbf{H}_0$ and $\|\cdot\|$ is the Frobenius norm induced from the inner product in \mathbb{S}^n . It is clear that the problem (12) is equivalent to

$$\text{maximize } \{y_0 \mid g_\lambda(y_0) = 0\}.$$

Thus we can solve (12) by the following simple bisection method.

Algorithm A: A bisection method for solving (12).

Choose a sufficiently large $\lambda > 0$, and a small tolerance $\varepsilon > 0$. Set $\mathbf{Y}_1^0 = \mathbf{0}$. Suppose that an interval $[a_0, b_0]$ has been determined such that $\eta^d(\lambda, \mathbb{K}) \in [a_0, b_0]$. Then perform the following steps at the k th iteration:

Step 1. Set $y_0^k = (a_{k-1} + b_{k-1})/2$.

Step 2. Compute $\Pi_{\mathbb{K}^*}(\mathbf{G}_\lambda(y_0^k)) = \mathbf{Y}_1^k + \mathbf{Y}_2^k$ with $\mathbf{Y}_i^k \in \mathbb{K}_i^*$, $i = 1, 2$, by Algorithm C using \mathbf{Y}_1^{k-1} as the starting point.

Step 3. If $g_\lambda(y_0^k) < \varepsilon \max\{1, |y_0^k|\}$, set $a_k = y_0^k$; else, set $b_k = y_0^k$.

Step 4. If $b_k - a_k \leq 5 \times 10^{-4} \max\{|a_k|, |b_k|\}$, stop, end.

In our numerical experiments, we choose $\varepsilon = 10^{-12}$ in Algorithm A.

To determine an interval $[a_0, b_0]$ which contains $\eta^d(\lambda, \mathbb{K})$, we can loosely solve (12) by Algorithm B described in Section 4.1 to produce an $\mathbf{X}_0 \in \mathbb{K}$. Assuming that $\mathbf{H}_0 \bullet \mathbf{X}_0 \neq 0$, then we can generate a feasible point $\hat{\mathbf{X}}$ for (11) by setting $\hat{\mathbf{X}} = \mathbf{X}_0 / (\mathbf{H}_0 \bullet \mathbf{X}_0)$. As a result, we have that

$$\eta^d(\lambda, \mathbb{K}) \leq \eta^p(\lambda, \mathbb{K}) \leq (\mathbf{Q}_0 + \lambda\mathbf{H}_1) \bullet \hat{\mathbf{X}} =: b_{\text{test}}.$$

The following cases are considered in order to determine the interval $[a_0, b_0]$:

(i) If $b_{\text{test}} \leq -1$, consider the set

$$\mathcal{J}_- = \{l \mid \kappa^l b_{\text{test}} \text{ is feasible for (12), } l \text{ is a positive integer}\}$$

where $\kappa > 1$ is a given constant, say 2. Let l^* be the smallest integer in \mathcal{J}_- . We can set $a_0 = \kappa^{l^*} b_{\text{test}}$ and $b_0 = \kappa^{l^*-1} b_{\text{test}}$. Numerically, we regard $\kappa^{-l} b_{\text{test}}$ as feasible if

$g_\lambda(\kappa^{-l}b_{\text{test}}) < \varepsilon \max\{1, |\kappa^{-l}b_{\text{test}}|\}$ holds.

(ii) If $b_{\text{test}} \geq 1$, consider the set

$$\mathcal{J}_+ = \{l \mid \kappa^{-l}b_{\text{test}} \text{ is feasible for (12), } l \text{ is a positive integer}\}.$$

If \mathcal{J}_+ is nonempty, let l^* be the smallest integer in \mathcal{J}_+ . We can set $a_0 = \kappa^{-l^*}b_{\text{test}}$ and $b_0 = \kappa^{1-l^*}b_{\text{test}}$. On the other hand, if \mathcal{J}_+ is empty, then we know that $\eta^d(\lambda, \mathbb{K}) \leq 0$. Thus if $y_0 = -1$ is feasible for (12), we can set $a_0 = -1$, $b_0 = 0$. Otherwise, we know that $\eta^d(\lambda, \mathbb{K}) < -1$, and hence we can set $b_{\text{test}} = -1$ and determine a_0, b_0 as in case (i).

(iii) If $b_{\text{test}} \in (-1, 1)$, we can set $b_0 = b_{\text{test}}$ and $a_0 = -1$ if $y_0 = -1$ is feasible for (12). Otherwise, we set $b_{\text{test}} = -1$ and determine a_0, b_0 as in case (i).

4.1 A proximal alternating direction multiplier method for solving (12)

Here we design a proximal alternating direction multiplier (PADM) method [14] for solving the following problem:

$$\text{minimize } \{-\mathbf{b}^T \mathbf{y} \mid \mathcal{A}^*(\mathbf{y}) + \mathbf{Z}_1 + \mathbf{Z}_2 = \mathbf{C}, \mathbf{Z}_1 \in \mathbb{S}_+^n, \mathbf{Z}_2 \in \mathbb{N}\}, \quad (14)$$

where $\mathbf{C} \in \mathbb{S}^n$, $\mathbf{b} \in \mathbb{R}^m$ are given data and $\mathcal{A} : \mathbb{S}^n \rightarrow \mathbb{R}^m$ is a given linear map. Note that (12) is a special case of (14) with $\mathcal{A}^*(y_0) = y_0 \mathbf{H}_0$, $\mathbf{b} = 1$, and $\mathbf{C} = \mathbf{Q}_0 + \lambda \mathbf{H}_1$.

Consider the following augmented Lagrangian function associated with (14):

$$\begin{aligned} L(\mathbf{y}, \mathbf{Z}_1, \mathbf{Z}_2; \mathbf{X}) &= -\mathbf{b}^T \mathbf{y} + \mathbf{X} \bullet (\mathcal{A}^* \mathbf{y} + \mathbf{Z}_1 + \mathbf{Z}_2 - \mathbf{C}) + \frac{\sigma}{2} \|\mathcal{A}^* \mathbf{y} + \mathbf{Z}_1 + \mathbf{Z}_2 - \mathbf{C}\|^2 \\ &= -\mathbf{b}^T \mathbf{y} + \frac{\sigma}{2} \|\mathcal{A}^* \mathbf{y} + \mathbf{Z}_1 + \mathbf{Z}_2 + \frac{1}{\sigma} \mathbf{X} - \mathbf{C}\|^2 - \frac{1}{2\sigma} \|\mathbf{X}\|^2 \end{aligned} \quad (15)$$

where $\mathbf{X} \in \mathbb{S}^n$, $\mathbf{y} \in \mathbb{R}^m$, $\mathbf{Z}_1 \in \mathbb{S}_+^n$, $\mathbf{Z}_2 \in \mathbb{N}$, and $\sigma > 0$ is the penalty parameter. Let \mathcal{T} be a given self-adjoint positive semidefinite linear operator defined on $\mathbb{S}^n \times \mathbb{S}^n$. The PADM method solves (14) by performing the following steps at the k th iteration:

$$\begin{aligned} \mathbf{y}^{k+1} &= \operatorname{argmin} \left\{ L(\mathbf{y}, \mathbf{Z}_1^k, \mathbf{Z}_2^k; \mathbf{X}^k) \mid \mathbf{y} \in \mathbb{R}^m \right\} \\ & \quad (\mathbf{Z}_1^{k+1}, \mathbf{Z}_2^{k+1}) \\ &= \operatorname{argmin} \left\{ \begin{array}{l} L(\mathbf{y}^{k+1}, \mathbf{Z}_1, \mathbf{Z}_2; \mathbf{X}^k) \\ + \frac{\sigma}{2} \begin{pmatrix} \mathbf{Z}_1 - \mathbf{Z}_1^k \\ \mathbf{Z}_2 - \mathbf{Z}_2^k \end{pmatrix} \bullet \mathcal{T} \begin{pmatrix} \mathbf{Z}_1 - \mathbf{Z}_1^k \\ \mathbf{Z}_2 - \mathbf{Z}_2^k \end{pmatrix} \end{array} \mid \mathbf{Z}_1 \in \mathbb{S}_+^n, \mathbf{Z}_2 \in \mathbb{N} \right\} \\ &= \operatorname{argmin} \left\{ \begin{array}{l} 2\mathbf{R}_d^k \bullet (\mathbf{Z}_1 - \mathbf{Z}_1^k + \mathbf{Z}_2 - \mathbf{Z}_2^k) \\ + \begin{pmatrix} \mathbf{Z}_1 - \mathbf{Z}_1^k \\ \mathbf{Z}_2 - \mathbf{Z}_2^k \end{pmatrix} \bullet \left(\begin{pmatrix} \mathcal{I} & \mathcal{I} \\ \mathcal{I} & \mathcal{I} \end{pmatrix} + \mathcal{T} \right) \begin{pmatrix} \mathbf{Z}_1 - \mathbf{Z}_1^k \\ \mathbf{Z}_2 - \mathbf{Z}_2^k \end{pmatrix} \end{array} \mid \mathbf{Z}_1 \in \mathbb{S}_+^n, \mathbf{Z}_2 \in \mathbb{N} \right\} \\ \mathbf{X}^{k+1} &= \mathbf{X}^k + \beta \sigma (\mathcal{A}^* \mathbf{y}^{k+1} + \mathbf{Z}_1^{k+1} + \mathbf{Z}_2^{k+1} - \mathbf{C}), \end{aligned}$$

where $\beta \in (0, \frac{1+\sqrt{5}}{2})$ is the step-length and $\mathbf{R}_d^k = \mathcal{A}^* \mathbf{y}^{k+1} + \mathbf{Z}_1^k + \mathbf{Z}_2^k + \sigma^{-1} \mathbf{X}^k - \mathbf{C}$.

By specifically choosing \mathcal{T} to be $\mathcal{T} = \begin{pmatrix} \mathcal{I} & -\mathcal{I} \\ -\mathcal{I} & \mathcal{I} \end{pmatrix}$, the computation of $\mathbf{Z}_1^{k+1}, \mathbf{Z}_2^{k+1}$ above then reduces to computing the projections onto \mathbb{S}_+^n and \mathbb{N} , respectively. As a result, we obtain the following efficient PADM method for solving (14). It can be shown that the PADM method converges; we refer the reader to [14] for the proof.

Algorithm B: A PADM method for solving (14).

Choose $\mathbf{Z}_1^0 = \mathbf{Z}_2^0 = \mathbf{X}^0 = \mathbf{0}$, and $\sigma > 0$, iterate the following steps:

Step 1. Compute \mathbf{y}^{k+1} by solving the following linear system of equations:

$$\mathcal{A} \mathcal{A}^* \mathbf{y} = \frac{1}{\sigma} (\mathbf{b} - \mathcal{A}(\mathbf{X}^k)) - \mathcal{A}(\mathbf{Z}_1^k + \mathbf{Z}_2^k - \mathbf{C})$$

Step 2. Compute

$$\mathbf{Z}_1^{k+1} = \operatorname{argmin} \left\{ \mathbf{R}_d^k \bullet (\mathbf{Z}_1 - \mathbf{Z}_1^k) + \|\mathbf{Z}_1 - \mathbf{Z}_1^k\|^2 \mid \mathbf{Z}_1 \in \mathbb{S}_+^n \right\} = \Pi_{\mathbb{S}_+^n} \left(\mathbf{Z}_1^k - \frac{1}{2} \mathbf{R}_d^k \right)$$

$$\mathbf{Z}_2^{k+1} = \operatorname{argmin} \left\{ \mathbf{R}_d^k \bullet (\mathbf{Z}_2 - \mathbf{Z}_2^k) + \|\mathbf{Z}_2 - \mathbf{Z}_2^k\|^2 \mid \mathbf{Z}_2 \in \mathbb{N} \right\} = \Pi_{\mathbb{N}} \left(\mathbf{Z}_2^k - \frac{1}{2} \mathbf{R}_d^k \right).$$

Step 3. Compute

$$\mathbf{X}^{k+1} = \mathbf{X}^k + \beta \sigma (\mathcal{A}^* \mathbf{y}^{k+1} + \mathbf{Z}_1^{k+1} + \mathbf{Z}_2^{k+1} - \mathbf{C}).$$

We can apply Algorithm B to (12) by taking $\mathbf{C} = \mathbf{Q}_0 + \lambda \mathbf{H}_1$, $\mathcal{A}(\mathbf{X}) = \mathbf{H}_0 \bullet \mathbf{X}$ and $\mathbf{b} = 1 \in \mathbb{R}$. Note that since the dual variable \mathbf{y} is one-dimensional, solving the linear system of equation in Step 1 is very easy.

4.2 Computing $\Pi_{\mathbb{K}}(\mathbf{G})$ and $\Pi_{\mathbb{K}^*}(-\mathbf{G})$

Observe that checking whether a given scalar y_0 is feasible for (12) amounts to checking whether $\mathbf{G}_\lambda(y_0) = \mathbf{Q}_0 + \lambda \mathbf{H}_1 - y_0 \mathbf{H}_0 \in \mathbb{K}^*$, i.e., whether $\Pi_{\mathbb{K}}(-\mathbf{G}_\lambda(y_0)) = \mathbf{0}$. To compute $\Pi_{\mathbb{K}}(\mathbf{G})$ for a given $\mathbf{G} \in \mathbb{S}^n$, we consider the following projection problem:

$$\operatorname{minimize} \left\{ \frac{1}{2} \|\mathbf{X} - \mathbf{G}\|^2 \mid \mathbf{X} \in \mathbb{K} \right\} \quad (16)$$

where the unique solution gives the projection $\Pi_{\mathbb{K}}(\mathbf{G})$ of \mathbf{G} onto \mathbb{K} . Note that for any $\mathbf{G} \in \mathbb{S}^n$, we have that $\mathbf{G} = \Pi_{\mathbb{K}}(\mathbf{G}) - \Pi_{\mathbb{K}^*}(-\mathbf{G})$ by the decomposition theorem of Moreau [20]. Using this equality, we can prove that $\mathbf{X} = \Pi_{\mathbb{K}}(\mathbf{G})$, $\mathbf{Y}_1 + \mathbf{Y}_2 = \Pi_{\mathbb{K}^*}(-\mathbf{G})$, $\mathbf{Y}_1 \in \mathbb{K}_1^*$

and $\mathbf{Y}_2 \in \mathbb{K}_2^*$ if and only if

$$\begin{aligned} \mathbf{X} - \mathbf{G} - \mathbf{Y}_1 - \mathbf{Y}_2 &= \mathbf{0}, \quad \mathbf{X} \bullet \mathbf{Y}_1 = 0, \quad \mathbf{X} \bullet \mathbf{Y}_2 = 0, \\ \mathbf{X} &\in \mathbb{K}_1, \quad \mathbf{X} \in \mathbb{K}_2, \quad \mathbf{Y}_1 \in \mathbb{K}_1^*, \quad \mathbf{Y}_2 \in \mathbb{K}_2^*, \end{aligned} \quad (17)$$

which can also be derived as the KKT condition for (16).

In Algorithm C described below, we will measure the accuracy of an approximation $(\hat{\mathbf{X}}, \hat{\mathbf{Y}}_1, \hat{\mathbf{Y}}_2)$ of $(\mathbf{X}, \mathbf{Y}_1, \mathbf{Y}_2)$ satisfying (17) by computing the following residual:

$$\delta_{\mathbb{K}} = \frac{1}{1 + \|\mathbf{G}\|} \max \left\{ \begin{array}{l} \|\hat{\mathbf{X}} - \mathbf{G} - \hat{\mathbf{Y}}_1 - \hat{\mathbf{Y}}_2\|, \frac{|\langle \hat{\mathbf{X}}, \hat{\mathbf{Y}}_1 \rangle|}{1 + \|\hat{\mathbf{Y}}_1\|}, \frac{|\langle \hat{\mathbf{X}}, \hat{\mathbf{Y}}_2 \rangle|}{1 + \|\hat{\mathbf{Y}}_2\|}, \\ \|\Pi_1^*(-\hat{\mathbf{X}})\|, \|\Pi_2^*(-\hat{\mathbf{X}})\|, \|\Pi_1(-\hat{\mathbf{Y}}_1)\|, \|\Pi_2(-\hat{\mathbf{Y}}_2)\| \end{array} \right\}. \quad (18)$$

It turns out that to solve (16) for a given $\mathbf{G} \in \mathbb{S}^n$, it is more efficient to consider the following problem for computing the projection of $-\mathbf{G}$ onto $\mathbb{K}^* = \mathbb{K}_1^* + \mathbb{K}_2^*$:

$$\text{minimize } \left\{ \text{minimize } \left\{ \frac{1}{2} \|\mathbf{G} + \mathbf{Y}_1 + \mathbf{Y}_2\|^2 \mid \mathbf{Y}_2 \in \mathbb{K}_2^* \right\} \mid \mathbf{Y}_1 \in \mathbb{K}_1^* \right\},$$

which can equivalently be formulated as follows:

$$\text{minimize } \left\{ f(\mathbf{Y}_1) := \frac{1}{2} \|\Pi_2(\mathbf{G} + \mathbf{Y}_1)\|^2 \mid \mathbf{Y}_1 \in \mathbb{K}_1^* \right\}. \quad (19)$$

It is easy to show that if \mathbf{Y}_1^* is an optimal solution of (19), then $\mathbf{X}^* = \Pi_2(\mathbf{G} + \mathbf{Y}_1^*)$ is an optimal solution of (16). It can be shown that $f(\cdot)$ is continuously differentiable on \mathbb{S}^n with $\nabla f(\mathbf{Y}_1) = \Pi_2(\mathbf{G} + \mathbf{Y}_1)$, and the gradient is Lipschitz continuous with modulus $L = 1$. However, note that $f(\cdot)$ is not twice continuously differentiable on \mathbb{S}^n .

The KKT conditions for (19) are given by:

$$\Pi_2(\mathbf{G} + \mathbf{Y}_1) - \mathbf{X} = \mathbf{0}, \quad \mathbf{Y}_1 \bullet \mathbf{X} = 0, \quad \mathbf{Y}_1 \in \mathbb{K}_1^*, \quad \mathbf{X} \in \mathbb{K}_1.$$

To solve the problem (19), we can either use a projected gradient method [5] or the accelerated proximal gradient (APG) method in [4]. We prefer the latter because of its superior iteration complexity. The details of the APG method is described next.

Algorithm C: An accelerated proximal gradient method for solving (19).

Choose $\bar{\mathbf{Y}}_1^0 = \mathbf{Y}_1^0 \in \mathbb{S}^n$ and a small tolerance $\epsilon > 0$; set $t_0 = 1$. Iterate the following steps:

Step 1. Compute $\nabla f(\bar{\mathbf{Y}}_1^k) = \Pi_2(\mathbf{G} + \bar{\mathbf{Y}}_1^k) = \max\{\mathbf{G} + \bar{\mathbf{Y}}_1^k, \mathbf{O}\}$.

Step 2. Set $\mathbf{H}^k = \bar{\mathbf{Y}}_1^k - L^{-1} \nabla f(\bar{\mathbf{Y}}_1^k)$. Compute $\mathbf{Y}_1^{k+1} = \Pi_1^*(\mathbf{H}^k)$.

Step 3. Set $t_{k+1} = \frac{1 + \sqrt{1 + 4t_k^2}}{2}$ and $\bar{\mathbf{Y}}_1^{k+1} = \mathbf{Y}_1^{k+1} + \frac{t_k - 1}{t_{k+1}} (\mathbf{Y}_1^{k+1} - \mathbf{Y}_1^k)$.

Step 4. Set $(\hat{\mathbf{X}}, \hat{\mathbf{Y}}_1, \hat{\mathbf{Y}}_2) = (\Pi_2(\mathbf{G} + \mathbf{Y}_1^{k+1}), \mathbf{Y}_1^{k+1}, \Pi_2^*(-\mathbf{G} - \mathbf{Y}_1^{k+1}))$. Compute the residual $\delta_{\mathbb{K}}$ defined in (18).

Step 5. If $\delta_{\mathbb{K}} \leq \epsilon$, stop, end.

We note that Algorithm C is called in Algorithm A with $\mathbf{G} = -\mathbf{G}_\lambda(y_0^k)$ to determine whether y_0^k is feasible for (12). It is also used to check whether $\kappa^l b_{test}$ and $\kappa^{-l} b_{test}$ are feasible for (12). As Algorithm C generates approximations of $\Pi_{\mathbb{K}}(\mathbf{G})$ and $\Pi_{\mathbb{K}^*}(-\mathbf{G})$, it is important to pick a small tolerance ϵ (10^{-12} in our numerical experiments) in the algorithm in order to determine the feasibility of y_0^k unambiguously.

We close this section with the following iteration complexity result for Algorithm C.

Theorem 4.1. *Let $\{\mathbf{Y}_1^k\}_{k=0}^\infty$ be generated by Algorithm C. Then for any $k \geq 1$, we have*

$$0 \leq f(\mathbf{Y}_1^k) - f(\mathbf{Y}_1^*) \leq \frac{2\|\mathbf{Y}_1^k - \mathbf{Y}_1^0\|^2}{(k+1)^2}.$$

Proof. It follows from [4, Theorem 4.1] by noting that $\nabla f(\cdot)$ is Lipschitz continuous with modulus $L = 1$. \square

5 Numerical experiments

To apply the numerical methods presented in the previous section, we take $\mathbb{K} = \mathbb{S}_+^n \cap \mathbb{N}$. We demonstrate the efficiency and effectiveness of solving the Lagrangian-DNN relaxation (12) using Algorithm A by comparing with the DNN relaxation (7) of (6) in terms of the quality of lower bounds and CPU time.

The test problems include the binary integer quadratic problems, the quadratic multiple knapsack problems, and the maximum stable set problems, and the quadratic assignment problems. All the experiments were performed in MATLAB on a Dell Precision T3500 Desktop with Intel Xeon quad-core CPU (2.80GHZ) and 24 GB memory.

5.1 Binary integer quadratic problems

The binary integer quadratic (BIQ) problem is described as

$$v^* = \min \{ \mathbf{x}^T \mathbf{Q} \mathbf{x} \mid \mathbf{x} \in \{0, 1\}^m \}, \quad (20)$$

where \mathbf{Q} is an $m \times m$ symmetric matrix (not necessarily positive semidefinite). By [8], a natural DNN relaxation of the problem (20) is given by:

$$v^{(0)} := \min \left\{ \mathbf{Q} \bullet \mathbf{X} \mid \begin{array}{l} \text{diag}(\mathbf{X}) - \mathbf{x} = 0, \\ \begin{bmatrix} 1 & \mathbf{x}^T \\ \mathbf{x} & \mathbf{X} \end{bmatrix} \in \mathbb{S}_+^{m+1} \cap \mathbb{N}^{m+1} \end{array} \right\}. \quad (21)$$

Note that by introducing slack variables, $v_i = 1 - x_i$, $i = 1, \dots, m$, (20) can be reformulated in the form (1) as follows:

$$v^* = \min \{ \mathbf{x}^T \mathbf{Q} \mathbf{x} \mid \mathbf{x} + \mathbf{v} = \mathbf{e}, \mathbf{x} \circ \mathbf{v} = 0, \mathbf{x} \geq 0, \mathbf{v} \geq 0 \}. \quad (22)$$

where \circ denotes the operation of performing elementwise multiplication. Observe that the direct DNN relaxation of the form (10) for the problem (22) is given by

$$v^{(1)} := \min \left\{ Q \bullet X \left| \begin{array}{l} \text{diag}(\mathbf{X}) - \mathbf{x} = 0, \text{diag}(\mathbf{V}) - \mathbf{v} = 0, \text{diag}(\mathbf{W}) = 0 \\ \mathbf{x} + \mathbf{v} = \mathbf{e}, \begin{bmatrix} 1 & \mathbf{x}^T & \mathbf{v}^T \\ \mathbf{x} & \mathbf{X} & \mathbf{W}^T \\ \mathbf{v} & \mathbf{W} & \mathbf{V} \end{bmatrix} \in \mathbb{S}_+^{2m+1} \cap \mathbb{N}^{2m+1} \end{array} \right. \right\}. \quad (23)$$

We can consider the relaxation of the form (7) for (22) with $\mathbb{K} = \mathbb{S}_+^{2m+1} \cap \mathbb{N}^{2m+1}$, and the subsequent Lagrangian-DNN relaxation (11) and its dual (12) to generate a lower bound $v^{(2)}$ for (20).

Table 5.1 presents the numerical results we obtain for solving (21), the direct DNN relaxation (23), and the Lagrangian-DNN relaxation of (22). The test problems are the Billionnet-Elloumi instances from BIQMAC library [6]. The lower bounds $v_{\text{LB}}^{(0)}$ obtained from (21) are the current state-of-the-art, and they are computed by the advanced large scale SDP solver, SDPNAL, originally developed in [32]. The stopping tolerance for SDPNAL is set to 10^{-5} rather than the default value of 10^{-6} since the former is more efficient for the purpose of computing a lower bound for (21). We should mention that as the solver SDPNAL only produces an approximately primal-dual feasible solution for (21), the procedure in [18] is used to generate a valid lower bound $v_{\text{LB}}^{(0)}$ for $v^{(0)}$ based on the approximately feasible solution.

From the table, we may observe that the Lagrangian-DNN relaxation (12) of (22) can produce much stronger lower bounds $v_{\text{LB}}^{(2)}$ than the bounds $v_{\text{LB}}^{(0)}$ generated from the standard DNN relaxation problem (21), while the CPU times taken to compute the bounds $v_{\text{LB}}^{(2)}$ by Algorithm A are at most 2.5 times that taken to compute $v_{\text{LB}}^{(0)}$ by SDPNAL. The fact that more time is needed to compute $v_{\text{LB}}^{(2)}$ should not come as a surprise since the matrix variable involved has dimension $2m + 1$ whereas the corresponding variable for $v_{\text{LB}}^{(0)}$ has dimension $m + 1$. But we have seen that the formulation (22) is able to produce much stronger lower bound than that produced by the standard DNN relaxation (21) of (20).

Next we test the strength of the Lagrangian-DNN relaxation (12) of (22) by comparing the lower bound $v_{\text{LB}}^{(2)}$ with the lower bound $v_{\text{LB}}^{(1)}$ produced by the direct DNN relaxation (23) of (22). Notice that the bounds $v_{\text{LB}}^{(2)}$ are tighter than $v_{\text{LB}}^{(1)}$ for almost all test problems, except for bqp100-3 and bqp100-4 where the differences between $v_{\text{LB}}^{(2)}$ and $v_{\text{LB}}^{(1)}$ are very small. The bounds $v_{\text{LB}}^{(1)}$ are now much closer to $v_{\text{LB}}^{(2)}$ compared to the differences between $v_{\text{LB}}^{(0)}$ and $v_{\text{LB}}^{(2)}$, however, computing $v_{\text{LB}}^{(1)}$ takes much longer time than $v_{\text{LB}}^{(2)}$ or $v_{\text{LB}}^{(0)}$. Thus we see that solving the Lagrangian-DNN relaxation of (22) by our proposed Algorithm A can generate the tight lower bounds $v_{\text{LB}}^{(2)}$ efficiently.

Table 1: BIQ problems with $m = 100$ and $m = 250$: $\lambda = 10^6 \|\overline{\mathbf{Q}}\| / \|\mathbf{H}_1\|$. The values $v_{\text{LB}}^{(0)}$ are computed based on (21) using the SDPNAL algorithm in [32]. Similarly, the values $v_{\text{LB}}^{(1)}$ in (23) are computed using SDPNAL.

problem	optimal value v^*	$v_{\text{LB}}^{(0)}$ (SDPNAL)	time	$v_{\text{LB}}^{(1)}$ (SDPNAL)	time	$v_{\text{LB}}^{(2)}$ (Algo. A)	time
bqp100-1	-7.97000000e3	-8.380978995e3	27	-8.071032266e3	102	-8.04687500e3	29
bqp100-2	-1.10360000e4	-1.148973750e4	28	-1.108631912e4	77	-1.10449219e4	26
bqp100-3	-1.27230000e4	-1.315374085e4	36	-1.272419250e4	148	-1.27246094e4	29
bqp100-4	-1.03680000e4	-1.073250841e4	33	-1.036927798e4	147	-1.03710937e4	28
bqp100-5	-9.08300000e3	-9.487049678e3	41	-9.115484423e3	64	-9.08935547e3	30
bqp250-1	-4.56070000e4	-4.766681027e4	201	-4.637834459e4	621	-4.62695313e4	272
bqp250-2	-4.48100000e4	-4.722281064e4	202	-4.573238489e4	720	-4.56054688e4	377
bqp250-3	-4.90370000e4	-5.108018909e4	167	-4.959841323e4	685	-4.94921875e4	420
bqp250-4	-4.12740000e4	-4.331281079e4	271	-4.213361046e4	778	-4.20507812e4	303
bqp250-5	-4.79610000e4	-5.000450078e4	259	-4.857234124e4	671	-4.84570313e4	398

5.2 Quadratic multiple knapsack problems

The problem under consideration is the following:

$$v^* := \min \{ \mathbf{x}^T \mathbf{Q} \mathbf{x} \mid \mathbf{A} \mathbf{x} + \mathbf{s} = \mathbf{b}, \mathbf{x} \in \{0, 1\}^m, \mathbf{s} \geq 0 \}, \quad (24)$$

where $\mathbf{A} \in \mathbb{R}^{q \times m}$ and $\mathbf{b} \in \mathbb{R}^q$ have their entries all being positive. The problem (24), studied in [9, 26], is a generalization of the binary quadratic single knapsack problem. In particular, the method in [9] solved the problem up to $q = 5$ and $m = 50$. In our numerical experiments, we set $q = 10$, $m = 100$, and use the same matrix \mathbf{Q} with $m = 100$ as in the BIQ problems in Section 5.1. The matrix \mathbf{A} is generated randomly with its elements independently drawn from the uniform distribution on the interval $[0, 10]$. The vector \mathbf{b} is set to $\mathbf{b} = m\mathbf{e}$.

By the formulation of Burer in [8], the natural DNN relaxation of (24) is given by

$$v^{(0)} := \min \left\{ \mathbf{Q} \bullet \mathbf{X} \left| \begin{array}{l} \mathbf{A} \mathbf{x} + \mathbf{s} = \mathbf{b}, \text{diag}(\mathbf{X}) - \mathbf{x} = 0 \\ \left(\begin{bmatrix} \mathbf{a}_i \\ \mathbf{e}_i \end{bmatrix} \begin{bmatrix} \mathbf{a}_i \\ \mathbf{e}_i \end{bmatrix}^T \right) \bullet \begin{bmatrix} \mathbf{X} & \mathbf{W}^T \\ \mathbf{W} & \mathbf{S} \end{bmatrix} = b_i^2 \ (i = 1, \dots, q) \\ \begin{bmatrix} 1 & \mathbf{x}^T & \mathbf{s}^T \\ \mathbf{x} & \mathbf{X} & \mathbf{W}^T \\ \mathbf{s} & \mathbf{W} & \mathbf{S} \end{bmatrix} \in \mathbb{S}^{m+q+1} \cap \mathbb{N}^{m+q+1} \end{array} \right. \right\} \quad (25)$$

where \mathbf{a}_i^T denotes the i th row of \mathbf{A} , and \mathbf{e}_i is the i th unit vector in \mathbb{R}^q . We can introduce slack variables, $v_i = 1 - x_i$, $i = 1, \dots, m$, just as for the BIQ problems in Section 5.1, to

reformulate (26) into the form (1) as follows:

$$v^* = \min \left\{ \mathbf{x}^T \mathbf{Q} \mathbf{x} \mid \begin{array}{l} \left[\begin{array}{ccc} \mathbf{I}_m & \mathbf{I}_m & 0 \\ \mathbf{A} & 0 & \mathbf{I}_q \end{array} \right] \begin{bmatrix} \mathbf{x} \\ \mathbf{v} \\ \mathbf{s} \end{bmatrix} = \begin{bmatrix} \mathbf{e} \\ \mathbf{b} \end{bmatrix}, \\ \mathbf{x} \circ \mathbf{v} = 0, \mathbf{x} \geq \mathbf{0}, \mathbf{v} \geq \mathbf{0}, \mathbf{s} \geq \mathbf{0} \end{array} \right\}. \quad (26)$$

In the numerical results presented in Table 5.2, we report the lower bound $v_{\text{LB}}^{(0)}$ computed by SDPNAL for (25); the lower bound $v_{\text{LB}}^{(1)}$ computed by SDPNAL for the DNN relaxation of (26) based on Burer's formulation in [8]; and the lower bound $v_{\text{LB}}^{(2)}$ computed by Algorithm A for the Lagrangian-DNN relaxation associated with (26).

As we can see from the numerical results that the lower bound $v_{\text{LB}}^{(0)}$ is much weaker than $v_{\text{LB}}^{(1)}$ and $v_{\text{LB}}^{(2)}$, thus we will mainly compare the bounds $v_{\text{LB}}^{(1)}$ and $v_{\text{LB}}^{(2)}$. Table 5.2 shows that the bound $v_{\text{LB}}^{(2)}$ based on the Lagrangian-DNN relaxation introduced in this paper can be computed much more efficiently than the lower bound $v_{\text{LB}}^{(1)}$. Furthermore, $v_{\text{LB}}^{(2)}$ is stronger than $v_{\text{LB}}^{(1)}$.

Table 2: Quadratic multiple knapsack problems: $\lambda = 10^6 \|\overline{\mathbf{Q}}\| / \|\mathbf{H}_1\|$. The values $v_{\text{LB}}^{(0)}$ are computed based on (25) using the SDPNAL algorithm in [32].

problem	$v_{\text{LB}}^{(0)}$ (SDPNAL)	time	$v_{\text{LB}}^{(1)}$ (SDPNAL)	time	$v_{\text{LB}}^{(2)}$ (Algo. A)	time
qmk100-1	-3.839274027e3	53	-3.696358669e3	185	-3.64746094e3	34
qmk100-2	-4.700674021e3	44	-4.558158548e3	145	-4.51171875e3	36
qmk100-3	-4.800266728e3	50	-4.679702850e3	177	-4.62890625e3	71
qmk100-4	-4.876050590e3	61	-4.732551889e3	182	-4.67651367e3	54
qmk100-5	-4.009051102e3	44	-3.883856215e3	172	-3.82690430e3	53
qmk250-1	-1.944972732e4	152	-1.924816440e4	900	-1.90136719e4	447
qmk250-2	-1.978813105e4	141	-1.960199989e4	1005	-1.92968750e4	460
qmk250-3	-2.024463770e4	130	-2.003825141e4	846	-1.97949219e4	441
qmk250-4	-1.878872520e4	154	-1.862331681e4	1041	-1.83935547e4	485
qmk250-4	-1.984494800e4	136	-1.967123749e4	1223	-1.94433594e4	441

5.3 Maximum stable set problems

For a graph G with m nodes and edge set \mathcal{E} , the stability number $\alpha(G)$ is the cardinality of a maximal stable set of G [10], and

$$-\alpha(G) := \min \{ -\mathbf{e}^T \mathbf{x} \mid x_i x_j = 0 \ ((i, j) \in \mathcal{E}), \mathbf{x} \in \{0, 1\}^m \} \quad (27)$$

$$\geq \min \left\{ (-\mathbf{e}\mathbf{e}^T) \bullet (\mathbf{z}\mathbf{z}^T) \mid \begin{array}{l} \sum_{i=1}^n z_i^2 = 1, z_i z_j = 0 \\ ((i, j) \in \mathcal{E}), \mathbf{z} \geq \mathbf{0} \end{array} \right\}. \quad (28)$$

Table 3: Maximum stable set problems with $n = 256, 512$ and 1024 : $\lambda = 10^6 \|\overline{\mathbf{Q}}\|/\|\mathbf{H}_1\|$. The values $v_{\text{LB}}^{(1)}$ are computed based on (30) using the SDPNAL algorithm in [32].

problem	optimal value v^*	$v_{\text{LB}}^{(1)}$ (SDPNAL)	time	$v_{\text{LB}}^{(2)}$ (Algo. A)	time
1dc.256	-30	-3.002325924e1	38	-3.00048828e1	15
1et.256	-50	-5.447886964e1	20	-5.44921875e1	28
1tc.256	-63	-6.325525918e1	22	-6.32812500e1	34
1zc.256	-36	-3.733740409e1	8	-3.73413086e1	10
1dc.512	-52	-5.272842750e1	160	-5.27099609e1	125
1et.512	-100	-1.035793061e2	136	-1.03613281e2	200
1tc.512	-110	-1.125746351e2	371	-1.12841797e2	305
1zc.512	-62	-6.800808859e1	87	-6.80175781e1	158
1dc.1024	-94	-9.557482297e1	2570	-9.55810547e1	1182
1et.1024	-171	-1.820965698e2	1032	-1.82128906e2	1398
1tc.1024	-196	-2.042445571e2	4443	-2.04589844e2	1811
1zc.1024	-112	-1.280028889e2	514	-1.28051758e2	233

Note that (28) is derived from (27) by setting $\mathbf{z} = \mathbf{x}/\sqrt{\mathbf{e}^T \mathbf{x}}$ for $0 \neq \mathbf{x} \in \{0, 1\}^m$.

Let \mathbf{E}_{ij} be the $m \times m$ symmetric matrix whose elements are all zeros, except the (i, j) and (j, i) elements which are equal to 1. By setting

$$\mathbf{Q}_0 = -\mathbf{e}\mathbf{e}^T, \quad \mathbf{H}_0 = \mathbf{I}, \quad \mathbf{H}_1 = \sum_{(ij) \in \mathcal{E}} \mathbf{E}_{ij} \quad (29)$$

it is readily shown that (28) is equivalent to the problem (8) with $\mathbb{K} = \mathbf{\Gamma}$. A well-known DNN relaxation of (28) is the following:

$$v^{(1)} = \min \{ \mathbf{Q}_0 \bullet \mathbf{X} \mid \mathbf{I} \bullet \mathbf{X} = 1, \mathbf{E}_{ij} \bullet \mathbf{X} = 0 \ ((i, j) \in \mathcal{E}), \mathbf{X} \in \mathbb{S}_+^n \cap \mathbb{N} \}. \quad (30)$$

Table 5.3 presents the comparison of the lower bounds and computation times for the problem (28). The test problems are graph instances (arising from coding theory) collected by Neil Sloane [27]. The lower bounds $v_{\text{LB}}^{(1)}$ we generated from (30) are the current state-of-the-art, and they are computed by the advanced large scale SDP solver, SDPNAL. The lower bounds $v_{\text{LB}}^{(2)}$ are computed for the problem (28) based on the Lagrangian-DNN relaxation (12) by Algorithm A. We can see that the bounds $v_{\text{LB}}^{(1)}$ and $v_{\text{LB}}^{(2)}$ are very close for all test problems, and the CPU times to compute $v_{\text{LB}}^{(1)}$ and $v_{\text{LB}}^{(2)}$ are comparable, except for the problems 1dc.1024, 1tc.1024, 1zc.1024. For these problems, computing $v_{\text{LB}}^{(1)}$ by SDPNAL takes more than twice time computing $v_{\text{LB}}^{(2)}$ by Algorithm A.

5.4 Quadratic assignment problems

We will identify a matrix $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_n] \in \mathbb{R}^{n \times n}$ with the n^2 -vector $\mathbf{x} = \text{vec}(\mathbf{X}) = [\mathbf{x}_1; \dots; \mathbf{x}_n]$. Given matrices $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{n \times n}$, the quadratic assignment problem is:

$$\begin{aligned} v^* &:= \min \left\{ \mathbf{X} \bullet \mathbf{A} \mathbf{X} \mathbf{B}^T \mid \begin{array}{l} \mathbf{e}^T \mathbf{X} \mathbf{e}_j = 1 = \mathbf{e}_j^T \mathbf{X} \mathbf{e} \\ (j = 1, \dots, n), \mathbf{X} \in \{0, 1\}^{n \times n} \end{array} \right\} \\ &= \min \left\{ \mathbf{x}^T (\mathbf{B} \otimes \mathbf{A}) \mathbf{x} \mid \begin{array}{l} (\mathbf{e}_j^T \otimes \mathbf{e}^T) \mathbf{x} = 1 = (\mathbf{e}^T \otimes \mathbf{e}_j^T) \mathbf{x} \\ (j = 1, \dots, n), \mathbf{x} \in \{0, 1\}^{n^2} \end{array} \right\}. \end{aligned} \quad (31)$$

Here $\mathbf{e} \in \mathbb{R}^n$ denotes the vector of ones, $\mathbf{e}_j \in \mathbb{R}^n$ the j th coordinate unit vector, and \otimes denotes the Kronecker product. We can express the above problem in the form of (1) by introducing an additional n vector variables $\mathbf{v}_i = \mathbf{e} - \mathbf{x}_i$ $i = 1, \dots, n$, but the resulting Lagrangian-DNN relaxation (12) of (1) will involve matrices with dimensions $2n^2 \times 2n^2$. From the computational point of view, such a doubling of matrix dimension is expensive, and our numerical experience also show that it is costly and difficult to solve (12) to high accuracy with such a formulation. For this reason, we will consider the following alternative reformulation of (31) introduced by Povh and Rendl in [23]:

$$\begin{aligned} v^* &:= \min \left\{ \mathbf{X} \bullet (\mathbf{A} \mathbf{X} \mathbf{B}^T) \mid \mathbf{X}^T \mathbf{X} = \mathbf{I} = \mathbf{X} \mathbf{X}^T, \mathbf{X} \geq 0 \right\} \quad (32) \\ &= \min \left\{ (\mathbf{B} \otimes \mathbf{A}) \bullet \mathbf{Y} \mid \begin{array}{l} \sum_{i=1}^n \mathbf{Y}^{ii} = \mathbf{I}, \mathbf{I} \bullet \mathbf{Y}^{ij} = \delta_{ij} \ (1 \leq i, j \leq n), \\ \mathbf{E} \bullet \mathbf{Y} = n^2, \mathbf{Y} \in \mathbb{C}^* \end{array} \right\}. \end{aligned} \quad (33)$$

Here \mathbf{Y}^{ij} corresponds to the (i, j) -block in the expansion of $\mathbf{x} \mathbf{x}^T$ as an $n^2 \times n^2$ matrix, and δ denotes Kronecker's delta. A natural DNN relaxation of (33) is following:

$$v^{(1)} := \min \left\{ (\mathbf{B} \otimes \mathbf{A}) \bullet \mathbf{Y} \mid \begin{array}{l} \sum_{i=1}^n \mathbf{Y}^{ii} = \mathbf{I}, \mathbf{I} \bullet \mathbf{Y}^{ij} = \delta_{ij} \ (1 \leq i, j \leq n), \\ \mathbf{E} \bullet \mathbf{Y} = n^2, \mathbf{Y} \in \mathbb{S}_+^{n^2} \cap \mathbb{N} \end{array} \right\}. \quad (34)$$

Using the fact that $\{\mathbf{X} \in \mathbb{R}^{n \times n} \mid \mathbf{X}^T \mathbf{X} = \mathbf{I}_n, \mathbf{X} \geq 0\}$ completely characterizes the set of $n \times n$ permutation matrices, we can show that (32) can equivalently be formulated as follows:

$$v^* := \min \left\{ \mathbf{X} \bullet (\mathbf{A} \mathbf{X} \mathbf{B}^T) \mid \begin{array}{l} \mathbf{e}^T \mathbf{X} \mathbf{e}_i = 1 = \mathbf{e}_i^T \mathbf{X} \mathbf{e} \ (i = 1, \dots, n), \\ X_{it} X_{jt} = 0 = X_{ti} X_{tj} \ (t = 1, \dots, n, i \neq j), \\ \mathbf{X} \geq \mathbf{O} \end{array} \right\}. \quad (35)$$

By considering the Lagrangian-DNN relaxation of (11) and its dual (12) for (35), we can generate a lower bound for (35), denoted as $v_{\text{LB}}^{(2)}$.

Table 5.4 presents the lower bounds and computation times corresponding to the relaxation problems (34) and the Lagrangian-DNN relaxation of (35) with $\mathbb{K} = \mathbb{S}_+^{n^2} \cap \mathbb{N}$.

Table 4: Quadratic assignment problems: $\lambda = 10^5 \|\overline{\mathbf{Q}}\| / \|\mathbf{H}_1\|$. The values $v_{\text{LB}}^{(1)}$ are computed based on (34) using the SDPNAL algorithm in [32]. The notation † means that the value is not known to be optimal.

problem	optimal value v^*	$v_{\text{LB}}^{(1)}$ (SDPNAL)	time	$v_{\text{LB}}^{(2)}$ (Algo. A)	time
bur26a	5.42667000e6	5.42508491e6	1485	5.42578125e6	261
bur26b	3.81785200e6	3.81569956e6	1200	3.81640625e6	212
bur26c	5.42679500e6	5.42409564e6	1853	5.42500000e6	237
bur26d	3.82122500e6	3.81843768e6	1483	3.81875000e6	156
bur26e	5.38687900e6	5.38585258e6	2241	5.38476563e6	187
chr15a	9.89600000e3	9.87953412e3	212	9.89453125e3	19
chr15b	7.99000000e3	7.98997388e3	160	7.98828125e3	21
chr15c	9.50400000e3	9.50397182e3	141	9.50390625e3	14
chr18a	1.10980000e4	1.10863272e4	281	1.10976563e4	43
chr18b	1.53400000e3	1.53351902e3	193	1.53281250e3	96
chr20a	2.19200000e3	2.19118820e3	578	2.19140625e3	78
chr20b	2.29800000e3	2.29783783e3	665	2.29785156e3	78
chr20c	1.41420000e4	1.41316846e4	714	1.41406250e4	92
chr22a	6.15600000e3	6.15574509e3	1290	6.15429688e3	90
chr22b	6.19400000e3	6.19362785e3	990	6.19335938e3	87
chr25a	3.79600000e3	3.79559572e3	1968	3.79589844e3	159
nug20	2.57000000e3	2.50185661e3	184	2.50500000e3	43
nug25	3.74400000e3	3.61940440e3	555	3.62500000e3	139
nug30	6.12400000e3	5.93960496e3	1209	5.94726563e3	306
tai30a	1.81814600e6†	1.70543609e6	1115	1.70625000e6	347
tai30b	6.37117113e8	5.94933253e8	2649	5.98242188e8	854

The test problems were downloaded from [24]. We observe that the bounds $v_{\text{LB}}^{(2)}$ are slightly closer to v^* for the problems `nug**`, `tai30*`, and are comparable for the other problems. However, the lower bounds $v_{\text{LB}}^{(2)}$ can be computed much faster by Algorithm A than $v_{\text{LB}}^{(1)}$ by SDPNAL.

5.5 Comparison of DNN relaxation (9) and Lagrangina-DNN relaxation (12)

To give the reader an idea on how difficult it is to solve the problem (9) with $\mathbb{K} = \mathbb{S}_+^n \cap \mathbb{N}$ accurately, we apply Algorithm B to solve the problem, and the numerical results is presented in Table 5.5. In the table, the quantities R_p and R_d are the relative primal and dual infeasibilities for the problems (8) and (9), respectively. As can be observed from the table, even after running Algorithm B for 10000 iterations, the dual feasibility of the final

iterate is still quite large except for the maximum stable set problems. As a result, the corresponding dual objective value does not provide a valid lower bound for (8). However, by using the computed dual variables $\hat{\mathbf{y}} = (\hat{y}_0, \hat{y}_1)$, we can attempt to generate a valid lower bound as follows. Let $\mathbf{G}(y_0) = \mathbf{Q}_0 - \hat{y}_1 \mathbf{H}_1 - y_0 \mathbf{H}_0$. Set

$$v_{\text{LB}}^{(3)} = \begin{cases} \max\{y_0^j := \hat{y}_0(1 + 0.01 \times j) \mid \mathbf{G}(y_0^j) \in \mathbb{K}^*, j = 0, 1, \dots\} & \text{if } \hat{y}_0 < 0 \\ \max\{y_0^j := \hat{y}_0(1 - 0.01 \times j) \mid \mathbf{G}(y_0^j) \in \mathbb{K}^*, j = 0, 1, \dots\} & \text{if } \hat{y}_0 > 0 \\ \max\{y_0^j := -0.01 \times j \mid \mathbf{G}(y_0^j) \in \mathbb{K}^*, j = 0, 1, \dots\} & \text{if } \hat{y}_0 = 0. \end{cases} \quad (36)$$

As we can observe from Table 5.5, it is generally difficult to solve (9) to high accuracy by Algorithm B or the solver SDPNAL, except for the maximum stable set problems. This is not surprising since the problems generally do not attain the optimal values by Lemma 3.3 (i). Moreover, the time taken to solve (9) and the subsequent generation of the valid lower bound $v_{\text{LB}}^{(3)}$ by Algorithm B or SDPNAL is generally much longer than that taken to compute $v_{\text{LB}}^{(2)}$ based on (12) by Algorithm A. Worse still, the lower bound $v_{\text{LB}}^{(3)}$ generated is also inferior to $v_{\text{LB}}^{(2)}$, except for the maximum stable set problems.

Table 5: Performance of PADM (Algorithm B with a maximum of 10000 iterations) and SDPNAL in solving (9) with $\mathbb{K} = \mathbb{S}_+^n \cap \mathbb{N}$.

problem	$v_{\text{LB}}^{(2)}$ (Algo. A)	time	$v_{\text{LB}}^{(3)}$ (PADM)	time iter	$\max\{R_p, R_d\}$	$v_{\text{LB}}^{(3)}$ (SDPNAL)	time	$\max\{R_p, R_d\}$	
bqp100-1	-8.04687500e3	29	-8.21461716e3	100	10000	1.62e-3	-8.14239100e3	218	5.08e-4
bqp100-2	-1.10449219e4	26	-1.12193981e4	102	10000	1.82e-3	-1.11888727e4	367	4.61e-4
bqp100-3	-1.27246094e4	29	-1.29551863e4	103	10000	1.80e-3	-1.29000349e4	170	1.45e-4
bqp100-4	-1.03710937e4	28	-1.05647288e4	114	10000	1.84e-3	-1.29000349e4	143	7.04e-5
bqp100-5	-9.08935547e3	30	-9.23988363e3	104	10000	1.81e-3	-9.20423625e3	236	8.51e-5
1dc.256	-3.00048828e1	15	-3.03000421e1	158	10000	1.57e-7	-3.02999610e1	22	8.60e-6
1et.256	-5.44921875e1	28	-5.50096590e1	125	10000	1.51e-7	-5.50124260e1	44	9.93e-6
1tc.256	-6.32812500e1	34	-6.38727706e1	127	10000	6.44e-7	-6.38726877e1	40	2.19e-6
1zc.256	-3.73413086e1	10	-3.77066666e1	23	2064	1.75e-9	-3.77062792e1	10	8.98e-6
chr15a	9.89453125e3	19	9.44046443e3	129	10000	1.53e-2	9.78302925e3	386	6.52e-4
chr15b	7.98828125e3	21	7.51407756e3	119	10000	2.19e-2	7.89900041e3	244	2.30e-4
chr15c	9.50390625e3	14	9.03523147e3	113	10000	2.40e-2	9.39679398e3	136	2.84e-4

6 Concluding remarks

We have presented a theoretical framework for the Lagrangian-conic relaxation of the QOP (1) and a computational method for the Lagrangian-DNN relaxation to improve the effectiveness and efficiency of obtaining the lower bounds for (1).

The theoretical results on the equivalence between the optimal value of the DNN relaxation of (1) and that of the Lagrangian-DNN relaxation shown in Section 3 provide the theoretical support for the tight lower bounds obtained numerically in Section 5. The computational efficiency of the proposed method has been achieved by the simple

bisection method combined with the efficient first order methods, the proximal alternating direction multiplier and the accelerated proximal gradient methods.

As shown in Section 5, the proposed method can compute the lower bounds of improved quality for the test problems efficiently. Specifically, the size of the quadratic multiple knapsack problems are larger than the ones solved in [9], and the quadratic assignment problem, known to be very hard problem to solve, could be handled very efficiently.

The approach proposed in this paper for the QOP (1) can be further extended to solve a general class of polynomial optimization problems (POPs). Currently, the numerical methods based on SDPs for POPs suffer from the numerical inefficiency caused by high degrees and large dimensions of POPs. Some of these difficulties can be dealt with the proposed idea in this paper. See also [3]. We intend to work on this project in the future.

References

- [1] N. Arima, S. Kim and M. Kojima. A quadratically constrained quadratic optimization model for completely positive cone programming. Research Report B-468, Department of Mathematical and Computing Sciences, Tokyo Institute of Technology, Oh-Okayama, Meguro-ku, Tokyo 152-8552, September, 2012. To appear in *SIAM J. Optim.*.
- [2] N. Arima, S. Kim and M. Kojima. Simplified Copositive and Lagrangian Relaxations for Linearly Constrained Quadratic Optimization Problems in Continuous and Binary Variables. Research Report B-469, Department of Mathematical and Computing Sciences, Tokyo Institute of Technology, Oh-Okayama, Meguro-ku, Tokyo 152-8552, October, 2012.
- [3] N. Arima, S. Kim and M. Kojima. Extension of completely positive cone relaxation to polynomial optimization, Research Report B-471, Department of Mathematical and Computing Sciences, Tokyo Institute of Technology, Oh-Okayama, Meguro-ku, Tokyo 152-8552, February, 2013.
- [4] A. Beck, and M. Teboulle. A fast iterative shrinkage-thresholding algorithm for linear inverse problems. *SIAM J. Imaging Sciences*, 2,183–202, 2009.
- [5] D. P. Bertsekas. *Nonlinear Programming*. Belmont MA, Athena Scientific, second edition, 1999.
- [6] BIQMAC Library, available at <http://www.biqmac.uni-klu.ac.at/biqmaclib.html>.
- [7] B. Borchers. CSDP, a C library for semidefinite programming. *Optim. Methods Softw.*, 11, 613–623, 1999.
- [8] S. Burer. On the copositive representation of binary and continuous non-convex quadratic programs. *Math. Program.*, 120, 479–495, 2009.

- [9] S. Burer. Optimizing a polyhedral-semidefinite relaxation of completely positive programs. *Math. Program. Comp.*, 2, 1–19, 2010.
- [10] E. de Klerk and D. V. Pasechnik, Approximation of the stability number of a graph via copositive programming, *SIAM J. Optim.*, 12, 875–892, 2002.
- [11] P. J. C. Dickinson, G. Eichfelder and J. Povh. Erratum to: “On the set-semidefinite representation of nonconvex quadratic programs over arbitrary feasible sets” [Optim. Letters, 2012], Optimization Online, http://www.optimization-online.org/DB_HTML/2012/09/3598.html, September 2012.
- [12] M. Dür and G. Still. Interior points of the completely positive cone. *Electron. J. Linear Al.*, 17, 28–33, 2008.
- [13] G. Eichfelder and J. Povh. On the set-semidefinite representation of nonconvex quadratic programs over arbitrary feasible sets. *Optim. Lett.*, DOI 10.1007/s11590-012-0450-3, 2012.
- [14] M. Fazel, Ti. K. Pong, D. Sun, and P. Tseng, Hankel matrix rank minimization with applications to system identification and realization. *SIAM J. Matrix Anal. A.*, 34, 946–977, 2013.
- [15] K. Fujisawa, M. Fukuda, K. Kobayashi, M. Kojima, K. Nakata, M. Nakata and M. Yamashita (2008) SDPA (SemiDefinite Programming Algorithm) user’s manual — Version 7.05. Research Report B-448, Dept. of Mathematical and Computing Sciences, Tokyo Institute of Technology, Oh-Okayama, Meguro, Tokyo 152-8552, Japan.
- [16] D. Gabay and B. Mercier, A dual algorithm for the solution of nonlinear variational problems via finite element approximations, *Comput. Math. Appl.*, 2, 17–40, 1976.
- [17] D. Ge and Y. Ye. On doubly positive semidefinite programming relaxations. Optimization Online, http://www.optimization-online.org/DB_HTML/2010/08/2709.html, August 2010.
- [18] C. Jansson and C. Keil. Rigorous error bounds for the optimal value in semidefinite programming. *SIAM J. Numer. Anal.*, 46, 188–200, 2007.
- [19] R. Glowinski and A. Marroco, *Sur l’approximation, par éléments finis d’ordre un, et la résolution, par pénalisation-dualité, d’une classe de problèmes de Dirichlet non linéaires. Revue Francaise d’Automatique, Informatique et Recherche Opérationnelle*, 9(R-2), pp. 41–76, 1975.
- [20] J. J. Moreau. Décomposition orthogonale d’un espace hilbertien selon deux cônes mutuellement polaires. *C. R. Acad. Sci.*, 255, 238–240, 1962.

- [21] K. G. Murty, S. N. Kabadi. Some NP-complete problems in quadratic and non-linear programming. *Math. Program.*, 39, 117–129, 1987.
- [22] Y. E. Nesterov and A. Nemirovskii, *Interior Point Methods for Convex Programming*. SIAM, Philadelphia, 1994.
- [23] J. Povh, and F. Rendl. Copositive and semidefinite relaxations of the quadratic assignment problem. *Discrete Optim.*, 6, 231–24, 2009.
- [24] Quadratic assignment problems, available at <http://www.seas.upenn.edu/qaplib>.
- [25] F. Rendl, G. Rinaldi, A. Wiegele. Solving max-cut to optimality by intersecting semidefinite and polyhedral relaxations. *Math. Program.* , 121, 307–335, 2010.
- [26] T. Sarac, A. Sipahioglu. A genetic algorithm for the quadratic multiple knapsack problem. In: *Advances in Brain, Vision, and Artificial Intelligence*, Vol. 4729 of Lecture Notes in Computer Science, pp. 490–498. Springer, Heidelberg, 2007.
- [27] N. Sloane. Challenge Problems: Independent sets in graphs. Available at <http://neilsloane.com/doc/graphs.html>
- [28] J. F. Strum. SeDuMi 1.02, a MATLAB toolbox for optimization over symmetric cones. *Optim. Method Softw.*, 11 & 12, 625–653, 1999.
- [29] K. Toh, M. J. Todd, R. H. Tütüntü. SDPT3 — a MATLAB software package for semidefinite programming. Dept. of Mathematics, National University of Singapore, Singapore, 1998.
- [30] A. W. Tucker. Dual systems of homogeneous linear relations. in Kuhn and Tucker eds., *Linear Inequalities and Related Systems*, Annals of Mathematics Studies Number 38, Princeton University Press, Princeton, N.J., 1956.
- [31] A. Yoshise and Y. Matsukawa. On optimization over the doubly nonnegative cone. *Proceedings of 2010 IEEE Multi-conference on Systems and Control*, 13–19, 2010.
- [32] X.Y. Zhao, D.F. Sun, and K.C. Toh. A Newton-CG augmented Lagrangian method for semidefinite programming. *SIAM J. Optim.*, 20, 1737–1765, 2010.